

ON AUTOMORPHISMS WITH NATURAL TANGENT ACTIONS ON HOMOGENEOUS PARABOLIC GEOMETRIES

JAN GREGOROVIČ AND LENKA ZALABOVÁ

ABSTRACT. We consider automorphisms of homogeneous parabolic geometries with a fixed point. Parabolic geometries carry the distinguished distributions and we study those automorphisms which enjoy natural actions on the distributions at the fixed points. We describe the sets of such automorphisms on homogeneous parabolic geometries in detail and classify, whether there are non-flat homogeneous parabolic geometries carrying such automorphisms. We present two general constructions of such geometries and we provide complete classifications for the types (G, P) of the parabolic geometries that have G simple and the automorphisms are of order 2.

INTRODUCTION

In this article, we will investigate automorphisms of Cartan geometries $(\pi : \mathcal{G} \rightarrow M, \omega)$ of type (G, P) over a connected smooth manifold that have fixed points and a natural tangent actions at that points. The crucial fact we will use is that the choice of a point $u \in \mathcal{G}$ in the fiber over the fixed point allows us to identify such automorphisms with elements of P . In particular, we can view u as a (higher order) frame of $T_{\pi(u)}M$, and the elements of P then prescribe the action of the automorphisms in this frame. The reduction to the first order frame of $T_{\pi(u)}M$ is then represented by the Lie group homomorphism $P \rightarrow Gl(\mathfrak{g}/\mathfrak{p})$ induced by the adjoint action. We would like to investigate automorphisms with natural tangent actions that do not depend on the choice of a preferred frame. In other words, we are interested in the elements that are mapped into the center of the image of the above Lie group homomorphism $P \rightarrow Gl(\mathfrak{g}/\mathfrak{p})$.

The affine symmetric spaces are our main motivation for the investigation of such automorphisms. Indeed, the unique non-trivial choice of a natural tangent action for the affine Cartan geometries is $-\text{id}_{\mathfrak{g}/\mathfrak{p}}$, i.e., the action of the geodesic symmetry. Thus the investigation of such automorphisms for Cartan geometries of an arbitrary type will give natural generalizations of symmetric spaces.

In this article, we will focus on regular parabolic geometries. These are Cartan geometries of type (G, P) , where G is semisimple and P is parabolic subgroup of G , and which are equivalent to a specific underlying geometric structures on M . The geometric data, which describe the underlying geometric structures depend on the type (G, P) . But in all cases, there is a distinguished maximally non-integrable distribution $T^{-1}M$ in TM , which is invariant under the action of all automorphisms. The remaining geometric data then restrict the possible frames of $T^{-1}M$ and admissible (partial) connections on $T^{-1}M$. Thus we investigate the

2010 *Mathematics Subject Classification.* 53C10; 53C30, 58J70.

Key words and phrases. parabolic geometries, homogeneous spaces, automorphisms with fixed points, harmonic curvature restrictions.

First author supported by the project CZ.1.07/2.3.00/20.0003 of the Operational Programme Education for Competitiveness of the Ministry of Education, Youth and Sports of the Czech Republic. Second author supported by the grant P201/11/P202 of the Czech Science Foundation (GAČR).

automorphisms with a fixed point $x \in M$ and a natural action on $T_x^{-1}M$ instead of the whole T_xM .

In particular, we investigate, which types of parabolic geometries admit non-flat homogeneous examples carrying automorphisms acting by a natural action on $T_x^{-1}M$. We will see that there are more possible natural actions than in the case of affine geometries.

In the first section, we recall basic facts on Cartan and parabolic geometries and their automorphisms. We distinguish the automorphisms according to their type Σ (see Definition 1). In particular, we consider the types Σ_J , which distinguish automorphisms with the same natural action J on $T_x^{-1}M$ (see Definition 2). Further, we distinguish the types of automorphisms, which generalize the geodesic symmetries of the affine symmetric spaces to parabolic geometries, and we call them generalized symmetries (see Definition 3). We also introduce all necessary notations and assumptions we use in this article.

We recall the structure of homogeneous parabolic geometries and the construction of them in the second section. We also describe the basic properties of automorphisms of arbitrary type Σ on homogeneous parabolic geometries and discuss how to compute all of them (see Theorem 2.4).

In the third section, we focus on the automorphisms of type Σ_J . We describe them in detail and classify all possible natural actions J on $T_x^{-1}M$ that can arise (see Proposition 3.2). The main result of this section is the Theorem 3.7, which describes the set of all automorphisms of type Σ_J .

The fourth section contains the results, which relate the existence of automorphisms of type Σ_J to the existence of generalized symmetries (see Theorem 4.1). This allows us to describe the curvature restrictions on these geometries in detail. Conversely, we give the restriction on the number of the generalized symmetries given by the existence of a non-trivial harmonic curvature (see Theorem 4.3).

We describe important classes of generalized symmetries in the fifth section. We show that these classes are in many cases closely related to symmetric spaces, and that the generalized symmetries in these cases are often coverings of the geodesic symmetries.

We describe two general constructions of non-trivial examples of homogeneous parabolic geometries with generalized symmetries in the sixth section. Moreover, we use the constructions to give several remarkable examples explicitly.

In the last section, we provide tables with the complete classifications for the types (G, P) of the parabolic geometries that have G simple, the automorphisms are of order 2 and that admit non-flat homogeneous parabolic geometries with these types of generalized symmetries.

1. STRUCTURE OF PARABOLIC GEOMETRIES AND THEIR AUTOMORPHISMS

Basic facts on Cartan geometries. Before we start to study the parabolic geometries, let us recall several general facts on Cartan geometries of an arbitrary type from [1, 12] we will use later. In particular, we discuss the automorphisms with a natural action on TM .

In the article, we will always consider a Cartan geometry $(\pi : \mathcal{G} \rightarrow M, \omega)$ of type (G, P) over a connected smooth manifold M with the curvature (function) $\kappa : \mathcal{G} \rightarrow \wedge^2(\mathfrak{g}/\mathfrak{p})^* \otimes \mathfrak{g}$. We denote by $\text{Aut}(\mathcal{G}, \omega)$ the automorphism group of the Cartan geometry $(\pi : \mathcal{G} \rightarrow M, \omega)$. Let us summarize here the crucial facts about $\text{Aut}(\mathcal{G}, \omega)$ we mentioned in the Introduction.

Fact The choice of $u \in \mathcal{G}$ provides the inclusion $\text{Aut}(\mathcal{G}, \omega) \rightarrow \mathcal{G}$ given by the evaluation of each automorphisms at u .

In particular, the transition $s \in P$ between the frames u and $\phi(u) = u \cdot s$ completely describes the action of the automorphisms ϕ with the fixed point $\pi(u)$. Since the Cartan connection ω identifies TM with $\mathcal{G} \times_{\underline{\text{Ad}}(P)} \mathfrak{g}/\mathfrak{p}$, where $\underline{\text{Ad}}$ is the truncated adjoint action on the quotient, the automorphism ϕ acts as

$$T_{\pi(u)}\phi \cdot \llbracket u, X + \mathfrak{p} \rrbracket = \llbracket u, \underline{\text{Ad}}_s(X + \mathfrak{p}) \rrbracket$$

on $T_{\pi(u)}M$.

The consequence of the first Fact is that we can identify the Lie subgroup in $\text{Aut}(\mathcal{G}, \omega)$ of all automorphisms ϕ with fixed point $\pi(u)$ with the Lie subgroup $A_u \subset P$ of all transitions between u and $\phi(u)$. Then $\underline{\text{Ad}}$ is a Lie group homomorphism of A_u to $Gl(\mathfrak{g}/\mathfrak{p})$, and the kernel of $\underline{\text{Ad}}$ describes the automorphisms with the same tangent action in the frame u . If we change the frame u to $u \cdot p$ for $p \in P$, then $p^{-1}sp$ is the element of A_{up} representing the action of automorphism ϕ . We will use the following notion for automorphisms with the distinguished action in some frame in the fiber over $\pi(u)$.

Definition 1. Let $(\pi : \mathcal{G} \rightarrow M, \omega)$ be a Cartan geometry of type (G, P) and Σ be a subset of P stable under the conjugation by elements of P . We say that an automorphism represented by $s \in A_u$ as above is an *automorphism of type Σ* at $\pi(u)$ if $s \in \Sigma$. For a fixed u , we call the elements of A_u automorphisms, too.

Since the set Σ is stable under the conjugation, the definition is independent of the actual choice of the frame u . On the other hand, the automorphisms of the same type Σ at $\pi(u)$ can have different tangent actions in the given frame u , because a priori $\underline{\text{Ad}}_s \neq \underline{\text{Ad}}_{p^{-1}sp}$. However, there are automorphisms, which share the same tangent action independently of the choice of the frame u . These are given by the elements of the center $Z(\underline{\text{Ad}}(P))$ of the image of $\underline{\text{Ad}}$. Such tangent actions are natural, and we can consider the types of automorphisms given by the preimage of elements in $Z(\underline{\text{Ad}}(P))$. Let us demonstrate this on affine Cartan geometries.

Example 1. Let us consider affine geometries, i.e., Cartan geometries of type $(\mathbb{R}^n \times Gl(n, \mathbb{R}), Gl(n, \mathbb{R}))$. Then \mathbb{R}^n can be identified with $\mathfrak{g}/\mathfrak{p}$ in a canonical way, and $\underline{\text{Ad}}$ is injective. In particular, the types of automorphisms with natural tangent actions correspond to $\Sigma = \{a \cdot \text{id}_{\mathbb{R}^n}\} \subset Z(Gl(n, \mathbb{R}))$ for $a \in \mathbb{R}$. Since the automorphisms of the Cartan geometry are exactly the affine transformations, they have to preserve the torsion T and the curvature R of the corresponding affine connection. It is clear that the automorphism of type $\{a \cdot \text{id}_{\mathbb{R}^n}\}$ at $\pi(u)$ act as $\frac{1}{a}$ on $T(\pi(u))$ and as $\frac{1}{a^2}$ on $R(\pi(u))$. Thus if we want to consider non-flat affine geometries with $a \cdot \text{id}_{\mathbb{R}^n} \in A_u$, then the only possible values of a are ± 1 . So the only non-trivial affine transformations we can consider are the geodesic symmetries corresponding to $a = -1$. The well known fact is the following:

Fact A homogeneous affine geometry such that the geodesic symmetry is at one (and thus at each) point of M a (global) affine transformation is a symmetric space.

Let us remark that for homogeneous affine geometries, the vanishing of the torsion is sufficient for $-\text{id}_{\mathbb{R}^n}$ to be an affine transformation.

Structure of the type (G, P) of parabolic geometries. Before we discuss the automorphisms of parabolic geometries with a natural action on the distinguished distribution $T^{-1}M$, let us make some assumption and recall the necessary technicalities on the structure of parabolic geometries. We follow here the book [1].

Let \mathfrak{g} be the Lie algebra (or the underlying real Lie algebra) of the real (complex) semisimple Lie group G . Then we will denote by α_i the simple restricted roots of \mathfrak{g} , and we consider their ordering according to [1, Appendix B]. There is the following crucial fact about the structure of the type (G, P) of parabolic geometries.

Fact There is a bijection between standard parabolic subalgebras of \mathfrak{g} and subsets Ξ of simple restricted roots (which are stable under the complex conjugation σ^*). Each parabolic subalgebra of \mathfrak{g} is conjugated to a standard one.

In this article, we will assume (without loss of generality) that the Lie algebra \mathfrak{p} of P is the standard parabolic subalgebra of \mathfrak{g} . Further, we will assume that the maximal normal subgroup of G contained in P is trivial, i.e., that the pair (G, P) is effective. For such parabolic geometries, id_G is the unique automorphism acting as id on M .

We use the notation P_{i_1, \dots, i_j} for the parabolic subgroup corresponding to the set $\Xi := \{\alpha_{i_1}, \dots, \alpha_{i_j}\}$. The set Ξ defines a functional ht_Ξ on the weight space of \mathfrak{g} given on simple restricted roots by $ht_\Xi(\alpha_i) = 1$ if $\alpha_i \in \Xi$ and $ht_\Xi(\alpha_i) = 0$ otherwise. We will denote by $\mu^\mathfrak{g}$ the highest root of \mathfrak{g} and define $k := ht_\Xi(\mu^\mathfrak{g})$. Then the structure of \mathfrak{g} is described as follows:

- Fact**
- (a) The functional ht_Ξ defines a $|k|$ -grading $\mathfrak{g}_{-k} \oplus \dots \oplus \mathfrak{g}_0 \oplus \dots \oplus \mathfrak{g}_k$ of \mathfrak{g} by $\mathfrak{g}_\alpha \subset \mathfrak{g}_{ht_\Xi(\alpha)}$ for the restricted root space \mathfrak{g}_α of the root α .
 - (b) There is an induced P -invariant filtration of \mathfrak{g} of the form $\mathfrak{g}^{-k} \supset \dots \supset \mathfrak{g}^0 = \mathfrak{p} \supset \dots \supset \mathfrak{g}^k$ given by $\mathfrak{g}^i := \mathfrak{g}_i \oplus \dots \oplus \mathfrak{g}_k$.
 - (c) The subgroup P is contained in the subgroup of filtration preserving elements of G , and contains its component of identity.

There always are distinguished subalgebras \mathfrak{g}_0 , $\mathfrak{p}_+ := \mathfrak{g}^1$ and $\mathfrak{g}_- = \mathfrak{g}_{-k} \oplus \dots \oplus \mathfrak{g}_{-1}$. The parabolic group P decomposes into a semidirect product of the Lie group G_0 with the Lie algebra \mathfrak{g}_0 , which consists of elements of P preserving the grading of \mathfrak{g} , and $\exp(\mathfrak{p}_+)$, where G_0 naturally acts by the adjoint action. Moreover, the grading induces a finer structure of P , and we can write each $p \in P$ uniquely as $p = g_0 \exp Z_1 \dots \exp Z_k$, where $g_0 \in G_0$ and $Z_i \in \mathfrak{g}_i$ for $i = 1, \dots, k$.

Automorphisms with a natural action on $T^{-1}M$. The consequence of the structure of the pair (G, P) is that all objects of the parabolic geometry $(\pi : \mathcal{G} \rightarrow M, \omega)$ of type (G, P) are filtered. In particular, there is a filtration $T^i M$ of TM which is induced by the filtration of $\mathfrak{g}/\mathfrak{p}$, and there always is a distinguished distribution $T^{-1}M$, the smallest piece of the filtration of TM , which is isomorphic to $\mathcal{G} \times_{\text{Ad}(P)} \mathfrak{g}^{-1}/\mathfrak{p}$. As we mentioned before, the distribution $T^{-1}M$ is the basic part of the geometric data describing the underlying geometric structure, so it is natural to study automorphisms with a fixed point $\pi(u)$ and a natural action on $T_{\pi(u)}^{-1}M$. In particular, the preimage of $Z(\text{Ad}|_{\mathfrak{g}^{-1}/\mathfrak{p}}(P))$ in $P = G_0 \rtimes \exp(\mathfrak{p}_+)$ is $Z(G_0) \rtimes \exp(\mathfrak{p}_+)$, because $\exp(\mathfrak{p}_+)$ acts trivially on $\mathfrak{g}^{-1}/\mathfrak{p}$.

Definition 2. Let $J \in Z(\text{Ad}|_{\mathfrak{g}^{-1}/\mathfrak{p}}(P))$. We denote by Σ_J the preimage of J in P , i.e.,

$$\Sigma_J = \{s \exp(Z) \in Z(G_0) \rtimes \exp(\mathfrak{p}_+) : \text{Ad}_s = J\}.$$

Thus we can fix an arbitrary frame u and use the Lie group A_u to study the Lie group of automorphisms fixing $\pi(u)$, because the results for the automorphisms of type Σ_J does not depend on the actual choice of u . In particular, this choice of u allows us to identify the Lie algebra of infinitesimal automorphisms $T_e \text{Aut}(\mathcal{G}, \omega)$ with the linear subspace $\text{inf}(u) \subset \mathfrak{g}$ given by the trivialization of $T_e \text{Aut}(\mathcal{G}, \omega) \subset T_u \mathcal{G} \cong \mathfrak{g}$. We note that the curvature $\kappa(u)$ measures the difference between the Lie algebra structure of \mathfrak{g} and $T_e \text{Aut}(\mathcal{G}, \omega)$. In particular, we will view $\text{inf}(u)$ as the Lie algebra of $\text{Aut}(\mathcal{G}, \omega)$, and denote by $\mathfrak{a}_u \subset \text{inf}(u)$ the Lie algebra of A_u .

Let us look on the action of automorphisms of type Σ_J on the neighborhood of the fixed point $\pi(u)$. If we consider the normal coordinates $\pi \circ \text{Fl}_1^{\omega^{-1}(X)}(u)$ for X in some neighborhood of 0 in \mathfrak{g}_- , then we can easily write down the action of

automorphisms $s \exp(Z) \in A_u \cap \Sigma_J$. Precisely, the automorphism $s \exp(Z)$ acts as

$$\pi \circ \mathrm{Fl}_1^{\omega^{-1}(X)}(u) \mapsto \pi \circ \mathrm{Fl}_1^{\omega^{-1}(X)}(us \exp(Z)).$$

Then s provides the linear change of coordinates of X in \mathfrak{g}_- , but the $\exp(\mathfrak{p}_+)$ -part changes the normal coordinate system itself. Thus apart the automorphisms of type Σ_J , there is another distinguished class of automorphisms with natural properties.

Definition 3. Let $(\pi : \mathcal{G} \rightarrow M, \omega)$ be a parabolic geometry of type (G, P) and $s \in Z(G_0)$. We say that automorphisms of type

$$\Sigma(s) := \{p^{-1}sp : p \in P\}$$

are s -symmetries. We say that automorphisms of type $\bigcup_{s \in Z(G_0)} \Sigma(s)$ are *generalized symmetries*.

We see immediately that $(\underline{\mathrm{Ad}}|_{\mathfrak{g}^{-1}/\mathfrak{p}})(\Sigma(s)) = \{\underline{\mathrm{Ad}}_s\}$ holds, i.e., s -symmetries have a natural action on $T^{-1}M$. Thus we will always write the elements of $Z(G_0)$ in the form s_J for $J = \underline{\mathrm{Ad}}_{s_J} \in Z(\underline{\mathrm{Ad}}|_{\mathfrak{g}^{-1}/\mathfrak{p}}(P))$. We will show in Proposition 3.1 that there is bijection between the actions J and elements s_J . Our main result of the fourth section states that on homogeneous parabolic geometries, the existence of automorphisms of type Σ_J induces the existence of the generalized symmetries with the same action on $T^{-1}M$.

Regularity and normality of parabolic geometries. Finally, let us add some further natural assumptions on the parabolic geometries we will investigate.

We will always assume that the parabolic geometry $(\pi : \mathcal{G} \rightarrow M, \omega)$ of type (G, P) is regular. In particular, the distribution $T^{-1}M$ is maximally non-integrable with the symbol \mathfrak{g}_- at all points of M . This means that $T^{-1}M$ generates the whole filtration $T^i M$ of TM , and it is reasonable to consider automorphisms with natural action just on $T^{-1}M$ and not on the whole TM . Indeed the action on $T^{-1}M$ of regular parabolic geometry induce the action on associated graded of TM .

Further, we will for simplicity assume that the parabolic geometry $(\pi : \mathcal{G} \rightarrow M, \omega)$ of type (G, P) is normal, i.e., that $\partial^* \kappa = 0$ holds for the curvature, where ∂^* is the Kostant codifferential. This allows to study the harmonic curvature (function) κ_H , which has its values in $\ker(\partial^*)/\mathrm{im}(\partial^*) \cong H^2(\mathfrak{g}_-, \mathfrak{g})$, instead of the whole curvature κ . In fact, κ_h is a basic invariant of regular, normal parabolic geometries, because there is a differential operator L such that $\kappa = L(\kappa_H)$, see [2].

The \mathfrak{g}_0 -module $H^2(\mathfrak{g}_-, \mathfrak{g})$ decomposes into a sum of isotypical components (\mathfrak{g}_0 -submodules) of \mathfrak{g}_0 -dominant weights, which have multiplicity 1. There are several possible ways, how to represent these isotypical components. We will use in this article the following ways:

- (1) Each component in $H^2(\mathfrak{g}_-, \mathfrak{g})$ is uniquely given by the ordered pair (i, j) such that the \mathfrak{g}_0 -dominant weight μ has the lowest weight vector $X^{-\mu}$ mapping $\mathfrak{g}_{-\alpha_i} \wedge \mathfrak{g}_{s_{\alpha_i}(-\alpha_j)}$ to $\mathfrak{g}_{s_{\alpha_i} s_{\alpha_j}(-\mu^{\mathfrak{g}})}$, where we denote by s_{α_i} the reflexion over the simple root α_i .
- (2) Each component in $H^2(\mathfrak{g}_-, \mathfrak{g})$ is uniquely given by the \mathfrak{g}_0 -dominant weight $\tilde{\mu}$ in $H^2(\mathfrak{p}_+, \mathfrak{g})$ given by the affine action of distinguished elements of the Hasse graph of the parabolic geometry (G, P) on the highest weight $\mu^{\mathfrak{g}}$.

The relation between the weights $\tilde{\mu}$ and μ is induced by the non-trivial isomorphism $H^2(\mathfrak{p}_+, \mathfrak{g}) \cong H^2(\mathfrak{g}_-, \mathfrak{g})$ and there are algorithms for the computation of both μ and $\tilde{\mu}$ for all components of the harmonic curvature, see [13].

Since $(\mathfrak{g}_-)^* = \mathfrak{p}_+$, we can equivalently view $X^{-\mu}$ as an element of $\mathfrak{p}_+ \wedge \mathfrak{p}_+ \otimes \mathfrak{g}$. Precisely, $X^{-\mu} = X^{\alpha_i} \wedge X^{s_{\alpha_i}(\alpha_j)} \otimes X^{s_{\alpha_i} s_{\alpha_j}(-\mu^{\mathfrak{g}})}$, where X^{α_i} is a root vector in the root space \mathfrak{g}_{α_i} .

2. HOMOGENEOUS PARABOLIC GEOMETRIES AND AUTOMORPHISMS OF TYPE Σ

Construction of homogeneous parabolic geometries. There is a general way how to construct all homogeneous parabolic geometries, see [1] or [7]. We recall the following definition.

Definition 4. Let (K, H) and (G, P) be types of Cartan geometries. Then the pair of maps (i, α) is called an *extension* of (K, H) to (G, P) if it satisfies the following conditions:

- (1) $i : H \rightarrow P$ is a Lie group homomorphism,
- (2) $\alpha : \mathfrak{k} \rightarrow \mathfrak{g}$ is a linear map extending $T_e i : \mathfrak{h} \rightarrow \mathfrak{p}$,
- (3) α induces an isomorphism $\mathfrak{k}/\mathfrak{h} \rightarrow \mathfrak{g}/\mathfrak{p}$ of vector spaces,
- (4) $\text{Ad}_{i(h)} \circ \alpha = \alpha \circ \text{Ad}_h$ holds for all $h \in H$.

Each extension (i, α) determines the *extension functor* \mathcal{F}_α from the category of Cartan geometries of type (K, H) to the category of Cartan geometries of type (G, P) . We have $\mathcal{F}_\alpha(\mathcal{B}) = \mathcal{B} \times_{i(H)} P$ for the Cartan geometry $(\mathcal{B} \rightarrow M, \omega)$ of type (K, H) , $\mathcal{F}_\alpha(\omega)$ is the unique Cartan connection, which pulls back to ω via the natural inclusion $\mathcal{B} \rightarrow \mathcal{F}_\alpha(\mathcal{B})$ and $\mathcal{F}_\alpha(\phi)$ for an automorphism $\phi \in \text{Aut}(\mathcal{B}, \omega)$ is the induced automorphism of the associated bundle $\mathcal{B} \times_{i(H)} P$.

There can be many extensions giving the same parabolic geometry. If we consider the natural inclusions $\text{Aut}(\mathcal{G}, \omega) \rightarrow \mathcal{G}$, then we obtain the inclusions $A_{u_0} \subset P$ and $\text{inf}(u_0) \subset \mathfrak{g}$ (which depend on the choice of u_0) which naturally satisfy all conditions of the extension for each choice of $u_0 \in \mathcal{G}$. We adopt the following terminology:

Definition 5. We say that the homogeneous parabolic geometry $(\mathcal{G} \rightarrow M, \omega)$ of type (G, P) is given by the extension (i, α) of (K, H) to (G, P) at $u_0 \in \mathcal{G}$, if there is an isomorphism between parabolic geometries $\mathcal{F}_\alpha(K \rightarrow K/H, \omega_K)$ and $(\mathcal{G} \rightarrow M, \omega)$ such that the class $[e, e] \in K \times_{i(H)} P$ corresponds to u_0 .

Let us point out that the kernel of the evaluation map at u_0 coincides with the kernel of i and consists of elements, which act as id on K/H . In particular, we will for simplicity assume that the pair (K, H) is effective, $K \subset \text{Aut}(\mathcal{G}, \omega)$ and i injective and we will omit writing it.

Automorphisms of type Σ on homogeneous parabolic geometries. Let $(\pi : \mathcal{G} \rightarrow M, \omega)$ be a homogeneous parabolic geometry of type (G, P) . Then we can consider the P -invariant subset $\bigcup_{u \in \mathcal{G}} (u, A_u)$ of the bundle $\mathcal{G} \times P$ with the right action of P given by $(u, q)p := (up, p^{-1}qp)$. The homogeneity allows to describe all automorphisms of the given type Σ at all points of M .

Proposition 2.1. *Let $(\pi : \mathcal{G} \rightarrow M, \omega)$ be a homogeneous parabolic geometry of type (G, P) and let $u_0 \in \mathcal{G}$. Then there is one-to-one correspondence between automorphisms of type Σ at any fixed point of M of the parabolic geometry $(\pi : \mathcal{G} \rightarrow M, \omega)$ and elements of $A_{u_0} \cap \Sigma$. In particular,*

$$\bigcup_{u \in \text{Aut}(\mathcal{G}, \omega)(u_0)} (u, A_u \cap \Sigma) \cong \text{Aut}(\mathcal{G}, \omega)(u_0) \times (A_{u_0} \cap \Sigma).$$

Proof. Clearly, $\phi^{-1}p\phi \in A_{u_0}$ holds for each $(\phi(u_0), p) \in \bigcup_{u \in \text{Aut}(\mathcal{G}, \omega)(u_0)} (u, A_u)$. Thus, since ϕ is invertible and unique for each point of $\text{Aut}(\mathcal{G}, \omega)(u_0)$, the second claim follows.

Since there always is $\phi \in \text{Aut}(\mathcal{G}, \omega)$ such that $\pi(\phi(u_0)) = x$ for any point $x \in M$, it follows that $\phi(A_{u_0} \cap \Sigma)\phi^{-1}$ is the set of automorphisms of type Σ at x , because the type of automorphism Σ is stable under conjugation. \square

Let $(\pi : \mathcal{G} \rightarrow M, \omega)$ be a homogeneous parabolic geometry of type (G, P) given by the extension (i, α) of effective (K, H) to (G, P) at $u_0 \in \mathcal{G}$. There is a natural

question, how to find all automorphisms of a chosen type Σ of this geometry, i.e., how to describe explicitly the set $A_{u_0} \cap \Sigma$?

The trouble is that K can be much smaller than the whole $\text{Aut}(\mathcal{G}, \omega)$. For example, we can view the flat model of (G, P) as an extension given by an inclusion of the maximal compact subgroup K into G . In particular, $H \cap \Sigma$ can be empty, even if $A_{u_0} \cap \Sigma$ is non-empty.

Lemma 2.2. *Let $(\pi : \mathcal{G} \rightarrow M, \omega)$ be a homogeneous parabolic geometry of type (G, P) given by the extension (i, α) of effective (K, H) to (G, P) at $u_0 \in \mathcal{G}$.*

- (1) *Let $h \in P$ be such that $\text{Ad}_h(\alpha(\mathfrak{k})) \subset \alpha(\mathfrak{k})$. Then $h \in A_{u_0}$ if and only if $h.\kappa(u_0) = \kappa(u_0)$ (for the action induced by Ad-action). In particular, such h induces an automorphism of \mathfrak{k} .*
- (2) *Denote*

$$\Sigma^\mathfrak{k} := \{h \in \Sigma : h.\kappa(u_0) = \kappa(u_0), \text{Ad}_h(\alpha(\mathfrak{k})) \subset \alpha(\mathfrak{k})\}.$$

Then there exists a pair (K', H') and an extension (α', i') of effective (K', H') to (G, P) giving the geometry $(\pi : \mathcal{G} \rightarrow M, \omega)$ at u_0 such that $i'(H') \cap \Sigma = \Sigma^\mathfrak{k}$.

- (3) *In particular, $\Sigma^\mathfrak{k}$ is the subset of $A_{u_0} \cap \Sigma$ consisting of all automorphisms, which preserve the K -orbits.*

Proof. If $h \in A_{u_0}$, then $\kappa(u_0) = \kappa(h(u_0)) = \kappa(u_0 h) = h.\kappa(u_0)$. Conversely, the following equality holds for the curvature:

$$\kappa(u_0)(X, Y) = h.\kappa(u_0)(X, Y) = \text{Ad}_h^{-1}([\alpha(X_h), \alpha(Y_h)] - \alpha([X_h, Y_h])),$$

where X_h, Y_h are arbitrary elements of $\text{inf}(u_0)$ congruent with $\text{Ad}_h(X), \text{Ad}_h(Y)$ modulo \mathfrak{p} , see [1, Section 1.5.16] for description of the curvature of homogeneous geometries. Since $\text{Ad}_h(\alpha(\mathfrak{k})) \subset \alpha(\mathfrak{k})$ and α is injective due to effectivity, we can take X_h to be the preimage of $\text{Ad}_h(\alpha(X))$ in \mathfrak{k} with respect to α for each $X \in \mathfrak{k}$. Thus $\text{Ad}_h([X, Y] - \alpha([X, Y])) = [\alpha(X_h), \alpha(Y_h)] - \alpha([X_h, Y_h])$. Comparing of the terms in the expression gives $[X, Y]_h = [X_h, Y_h]$ due to injectivity of α , i.e. h induces an automorphism of \mathfrak{k} .

If there is no $k \in H$ such that $\text{Ad}_k = \text{Ad}_h$, then we can form the semidirect products $(K \rtimes \{h^k : k \in \mathbb{Z}\}, H \rtimes \{h^k : k \in \mathbb{Z}\}) =: (K', H')$, and $i'(h', h^k) := i(h')h^k$ naturally extends to the extension (α, i') of (K', H') to (G, P) . Then the map $[[k, h], p] \mapsto [[k, hp]]$ is an isomorphism of parabolic geometries $\mathcal{F}_\alpha(K \rightarrow K/H, \omega_K)$ and $\mathcal{F}_{\alpha'}(K' \rightarrow K'/H', \omega_{K'})$. Thus $h \in A_{u_0}$ and the claim (1) follows.

The same construction can be done for the whole $\Sigma^\mathfrak{k}$ instead of $\{h^k : k \in \mathbb{Z}\}$, which implies the claim (2) directly. Then the claim (3) follows immediately. \square

In particular, we get the following consequence of the previous Lemma.

Corollary 2.3. *For $\theta \in \text{Aut}(\mathfrak{g})$ such that $\theta(\mathfrak{p}) \subset \mathfrak{p}$, there is the effective pair (G', P') such that $G/P \simeq G'/P'$ and $\theta = \text{Ad}_h$ for $h \in P'$.*

The distinguished case $\mathfrak{k} = \text{inf}(u_0)$ allows to describe the set $A_{u_0} \cap \Sigma$ explicitly. In particular, we will use the notation $\Sigma^{\text{inf}(u_0)}$ for the set of all automorphisms of type Σ at $\pi(u_0)$ rather than the notation $A_{u_0} \cap \Sigma$ in the rest of the article.

Theorem 2.4. *Let $(\pi : \mathcal{G} \rightarrow M, \omega)$ be a homogeneous parabolic geometry of type (G, P) . Then*

$$\Sigma^{\text{inf}(u_0)} = \{h \in \Sigma : h.\kappa(u_0) = \kappa(u_0), \text{Ad}_h(\text{inf}(u_0)) \subset \text{inf}(u_0)\} = A_{u_0} \cap \Sigma.$$

There is an algorithm for computation of the local infinitesimal automorphisms of homogeneous parabolic geometries, see [7]. Then $\text{inf}(u_0)$ consists of complete

local infinitesimal automorphisms, and the above Theorem shows how to compute automorphisms of type Σ , which are contained in $\Sigma^{\text{inf}(u_0)}$.

We will see in the next chapters, that the set $\Sigma^{\text{inf}(u_0)}$ can be computed more easily in the case of Σ_J .

3. HOMOGENEOUS PARABOLIC GEOMETRIES AND AUTOMORPHISMS OF TYPE Σ_J

Description of Σ_J and structure of $Z(G_0)$. We focus here on types Σ_J of automorphisms with a natural action on $T^{-1}M$. In particular, we describe the set Σ_J for $J \in Z(\underline{\text{Ad}}|_{\mathfrak{g}^{-1}/\mathfrak{p}}(P))$ in detail.

Proposition 3.1. *For the pair (G, P) , there is a bijection between actions $J \in Z(\underline{\text{Ad}}|_{\mathfrak{g}^{-1}/\mathfrak{p}}(P))$ and elements $s_J \in Z(G_0)$. In particular, for each $J \in Z(\underline{\text{Ad}}|_{\mathfrak{g}^{-1}/\mathfrak{p}}(P))$, there is a unique $s_J \in Z(G_0)$ such that*

$$\Sigma_J = \{s_J \exp(Z) : Z \in \mathfrak{p}_+\}$$

and $J = \text{Ad}_{s_J}|_{\mathfrak{g}^{-1}}$.

Proof. The action of $\underline{\text{Ad}}_s \in \underline{\text{Ad}}|_{\mathfrak{g}^{-1}/\mathfrak{p}}(\Sigma_J)$ on $\mathfrak{g}^{-1}/\mathfrak{p}$ coincides with the action of Ad_s on \mathfrak{g}_{-1} . Let $s'_J \in Z(G_0)$ be another element such that $\text{Ad}_{s'_J} = \text{Ad}_{s_J}$ on \mathfrak{g}_{-1} . Then $\text{Ad}_{s_J(s'_J)^{-1}} = \text{id}$ holds on \mathfrak{g}_- , because \mathfrak{g}_- is generated by \mathfrak{g}_{-1} . Since the geometry is effective, there is no ideal of \mathfrak{g} contained in \mathfrak{g}_0 . This means that $\text{Ad}_{s_J(s'_J)^{-1}} = \text{id}$ holds on the whole \mathfrak{g} thanks to the duality $\mathfrak{g}_-^* = \mathfrak{p}_+$ and the fact $\mathfrak{g}_0 \subseteq \mathfrak{g}_-^* \otimes \mathfrak{g}_-$. Thus the effectivity of (G, P) implies $s_J = s'_J$ and the claim follows. \square

Let us describe the elements of $Z(G_0)$ in the form of a functional on the space of weights.

Proposition 3.2. *Let \mathfrak{g} be a semisimple Lie algebra and \mathfrak{p} its parabolic subalgebra, which does not contain any simple ideal of \mathfrak{g} . Then the following facts hold:*

- (1) *For each pair (G, P) and each $s_J \in Z(G_0)$, there is a complex valued linear functional λ_J given on the space of weights of \mathfrak{g} given by*

$$\text{Ad}_{s_J}(X^\alpha) = e^{\lambda_J(\alpha)} X^\alpha$$

for the restricted root α . In particular, the following conditions are satisfied:

- (a) *If $X^\alpha \in \mathfrak{g}_0$, then $\lambda_J(\alpha) = 0$.*
 - (b) *If \mathfrak{g}_α is a real root space, then $\lambda_J(\alpha) \in \mathbb{R} \pmod{\pi i}$.*
 - (c) *If \mathfrak{g}_α is a complex root space, then $\lambda_J(\alpha)$ and $\lambda_J(\sigma^* \alpha)$ are complex conjugate.*
- (2) *Let λ_J be a complex valued linear functional on the space of weights of \mathfrak{g} satisfying (a)-(c). Then there is a pair (G, P) satisfying all our assumptions with $s_J \in Z(G_0)$ satisfying*

$$\text{Ad}_{s_J}(X^\alpha) = e^{\lambda_J(\alpha)} X^\alpha$$

for all restricted roots α .

- (3) *In particular, the unique eigenvalue of $s_J \in Z(G_0)$ on the component of the harmonic curvature (i, j) is $e^{\lambda_J(\mu)}$ for the corresponding weight μ , and if $X^{-\mu} = X^{\alpha_i} \wedge X^{s_{\alpha_i}(\alpha_j)} \otimes X^{s_{\alpha_i} s_{\alpha_j}(-\mu^{\mathfrak{g}})}$, then*

$$\text{Ad}_{s_J}(X^{-\mu}) = e^{\lambda_J(\alpha_i) + \lambda_J(s_{\alpha_i}(\alpha_j)) - \lambda_J(s_{\alpha_i} s_{\alpha_j}(\mu^{\mathfrak{g}}))} X^{-\mu}.$$

Proof. We get the claim (1) just by the restriction of the adjoint representation of G to G_0 and by the complete reducibility. Then the properties (a)–(c) trivially follows. Conversely, if the conditions (a)–(c) are satisfied, then the formula in the claim (2) induces an automorphism of \mathfrak{g} , and the claim (2) follows from the Corollary 2.3, because $\text{Ad}_{s_J}|_{\mathfrak{g}_0} = \text{id}$ implies $s_J \in Z(G_0)$. The claim (3) is then a direct consequence of the description of components of the harmonic curvature. \square

Clearly, the set Σ_J is a union of P -conjugacy classes of suitable elements of P . We show that there are distinguished representatives of the P -conjugacy classes.

Lemma 3.3. *There is a finite amount of $Z_i \in \mathfrak{p}_{fix}^{s_J}$ such that*

$$\Sigma_J = \Sigma(s_J) \cup \bigcup_i \Sigma(s_J \exp(Z_i)),$$

where

$$\mathfrak{p}_{fix}^{s_J} := \{Z \in \mathfrak{p}_+ : \text{Ad}_{s_J}(Z) = Z\}.$$

In particular, for $s_J \exp(Z) \in \Sigma_J$ there is $Y \in \mathfrak{p}_+$ such that $\exp(-Y)s_J \exp(Z) \exp(Y) \in s_J \exp(\mathfrak{p}_{fix}^{s_J})$ and

$$\Sigma(s_J) = \{s_J \exp(C(-\text{Ad}_{s_J}^{-1}(Y), Y)) : Y \in \mathfrak{p}_+\},$$

where C denotes the application of the Baker–Campbell–Hausdorff (BCH) formula on the elements, see [9, Theorem 4.29].

Proof. For $s_J \in Z(G_0)$, there always is only a finite amount of P -conjugacy classes

$$\Sigma(s_J \exp(Z)) = \{s_J \exp(-\text{Ad}_{s_J}^{-1}Y) \exp(\text{Ad}_{g_0}^{-1}Z) \exp(Y) : g_0 \in G_0, Y \in \mathfrak{p}_+\}$$

for $Z \in \mathfrak{p}_+$, because there always is a finite amount of $\text{Ad}(G_0)$ -orbits in \mathfrak{p}_+ .

Let us consider an element $\exp(Y) \in \exp(\mathfrak{p}_+)$. The conjugation of $s_J \exp(Z)$ by this element is of the form

$$\exp(-Y)s_J \exp(Z) \exp(Y) = s_J \exp(-\text{Ad}_{s_J}^{-1}Y) \exp(Z) \exp(Y).$$

Thus if $Z_f + Z'$ is the decomposition of Z into $\mathfrak{p}_{fix}^{s_J}$ and the remaining eigenspaces of Ad_{s_J} , then we can choose $Y := (\text{id} - \text{Ad}_{s_J}^{-1})^{-1}Z'$ to change the representative, because $(\text{id} - \text{Ad}_{s_J}^{-1})$ is invertible on eigenspaces of Ad_{s_J} for eigenvalues different from 1. Then the right hand side is of the form

$$s_J \exp(\text{Ad}_{s_J}^{-1}(\text{id} - \text{Ad}_{s_J}^{-1})^{-1}Z' + Z_f + Z' - (\text{id} - \text{Ad}_{s_J}^{-1})^{-1}Z' + \dots) = s_J \exp(Z_f + \dots),$$

where \dots are given by the BCH-formula. Since \dots are in the higher order part of the filtration than Z , we can repeat the computation for \dots . Then the result follows by induction, because the new terms appear in the part of the filtration containing $[Z_f, \dots]$ due to BCH-formula, and thus vanish after k steps of induction for $|k|$ -graded geometry due to nilpotency.

The last formula then corresponds to the case $Z = 0$. □

Jordan decomposition of automorphisms of type Σ_J . It is clear from the Proposition 3.2 that $s_J \in Z(G_0)$ are semisimple elements. This gives the following alternative conditions on s_J being an element of $\Sigma_J^\natural := (\Sigma_J)^\natural$.

Lemma 3.4. *Let $(\pi : \mathcal{G} \rightarrow M, \omega)$ be a homogeneous parabolic geometry given by the extension (i, α) of effective (K, H) to (G, P) at $u_0 \in \mathcal{G}$. Then the following claims hold:*

- (1) *The following facts are equivalent:*
 - (a) $\text{Ad}_{s_J} \cdot \kappa(u_0) = \kappa(u_0)$,
 - (b) *the components of the harmonic curvature such that $\lambda_J(\mu) \not\equiv 0 \pmod{2\pi i}$ for the corresponding weight μ vanish at u_0 .*
- (2) *The following facts are equivalent:*
 - (a) $\text{Ad}_{s_J}(\alpha(\mathfrak{k})) \subset \alpha(\mathfrak{k})$,
 - (b) $\alpha(\mathfrak{k})$ *splits into eigenspaces of Ad_{s_J} in \mathfrak{g} .*

In particular, if one of the points (a) or (b) is satisfied for both claims (1) and (2), then $s_J \in \Sigma_J^\natural$.

Proof. (1) If $\text{Ad}_{s_J} \cdot \kappa(u_0) = \kappa(u_0)$, then the same holds for κ_H and we know that $\lambda_J(\mu) = 0 \pmod{2\pi i}$ for the corresponding weight μ from Proposition 3.2. Conversely, the condition (b) implies $\text{Ad}_{s_J} \cdot \kappa_H(u_0) = \kappa_H(u_0 s_J) = \kappa_H(u_0)$ and the claim follows from the general theory, because the splitting operator giving the whole curvature is G_0 -equivariant, see [2, 1].

(2) The condition (b) clearly implies (a). Conversely, since s_J is a semisimple element, there is a basis of $\alpha(\mathfrak{k})$ consisting of eigenvectors of Ad_{s_J} and the claim follows.

In particular, $s_J \in \Sigma_J^{\mathfrak{k}}$ holds if the points (a) are satisfied for both claims (1) and (2) by the definition of $\Sigma_J^{\mathfrak{k}}$. \square

We know from the Lemma 3.3 that there is u_0 such that $s_J \exp(Z)$ is the Jordan decomposition of an element of Σ_J , i.e., s_J is semisimple element, $\exp(Z)$ unipotent element and $\text{Ad}_{s_J}(Z) = Z$. This has the following consequence.

Lemma 3.5. *Let $(\pi : \mathcal{G} \rightarrow M, \omega)$ be a homogeneous parabolic geometry of type (G, P) . Then there is $u_0 \in \mathcal{G}$ such that $s_J \in \Sigma_J^{\mathfrak{k}}$ for the extension (i, α) of effective (K, H) to (G, P) giving the geometry at u_0 with $\Sigma_J^{\mathfrak{k}} \neq \emptyset$.*

Proof. Let us choose the point u_0 such that $\text{Ad}_{s_J}(Z) = Z$ holds for the element $s_J \exp(Z) \in \Sigma_J^{\mathfrak{k}}$. Then since $s_J \exp(Z)$ is the Jordan decomposition, $\text{Ad}_{s_J}(\alpha(\mathfrak{k})) \subset \alpha(\mathfrak{k})$ holds, i.e., the condition (a) of the claim (2) of the Lemma 3.4 is satisfied. Moreover, since $s_J \exp(Z) \in \Sigma_J^{\mathfrak{k}}$, the condition (b) of the claim (1) of the Lemma 3.4 is satisfied, and the claim follows. \square

Let us adopt the following notation

$$\mathfrak{h}_+ := \mathfrak{h} \cap \mathfrak{p}_+.$$

Lemma 3.6. *Let $(\pi : \mathcal{G} \rightarrow M, \omega)$ be a homogeneous parabolic geometry given by the extension (i, α) of effective (K, H) to (G, P) at $u_0 \in \mathcal{G}$. Then*

$$H \cap \exp(\mathfrak{p}_+) = \Sigma_{\text{id}_{\mathfrak{g}}^{-1/\mathfrak{p}}}^{\mathfrak{k}} = \exp(\mathfrak{h}_+).$$

Proof. Let us write $X_i + \dots$ for the element $X \in \mathfrak{g}^i$ such that $0 \neq X_i \in \mathfrak{g}^i / \mathfrak{g}^{i+1} \simeq \mathfrak{g}_i$, and suppose $Z = Z_j + \dots$ for $\exp(Z) \in H \cap \exp(\mathfrak{p}_+)$. Then it follows from the general theory that there is an $\mathfrak{sl}(2)$ -triple (X_{-j}, A_0, Z_j) for $X_{-j} \in \mathfrak{g}_{-j}$ and $A_0 \in \mathfrak{g}_0$, see [8, Section X.3]. Since $j > 0$, there is $X \in \alpha(\mathfrak{k})$ such that $X = X_{-j} + \dots$ and we define $A := (\text{id}_{\mathfrak{k}} - \text{Ad}_{\exp(Z)})(X)$, i.e., $\alpha(A) = -[Z_j, X_{-j}] + \dots$ holds. Thus

$$(\text{id}_{\mathfrak{k}} - \text{Ad}_{\exp(Z)})^2(X) = [A_0, Z_j] + \dots = 2Z + Y \in \mathfrak{h}_+,$$

where $Y \in \mathfrak{p}_+$ is in the higher order part of the filtration than Z . In particular, if $Y = 0$, then $Z \in \mathfrak{h}$. Otherwise, $(\exp(Z))^2 \exp(-2Z - Y) = \exp(Y') \in H \cap \exp(\mathfrak{p}_+)$, where $Y' \in \mathfrak{p}_+$ is in the higher order part of the filtration than Z .

Thus the above computations can be seen as an induction step. If $Y' \in \mathfrak{h}$, then $\exp(2Z) = \exp(Y') \exp(2Z + Y)$, and thus $Z \in \mathfrak{h}_+$. Clearly, the filtration component of Y' grows by each step, and thus, we get $Y' = 0 \in \mathfrak{h}$ after k steps for $|k|$ -graded geometry and the claim follows by induction. \square

There is the following crucial consequence of the above Jordan decomposition from the Lemma 3.5, which in particular simplifies the computation of $\Sigma_J^{\text{inf}(u_0)}$.

Theorem 3.7. *Let $(\pi : \mathcal{G} \rightarrow M, \omega)$ be a homogeneous parabolic geometry of type (G, P) . Then there is $u_0 \in \mathcal{G}$ such that*

$$\Sigma_J^{\mathfrak{k}} = \{s_J \exp Z : Z \in \mathfrak{h}_+\}$$

for the extension (i, α) of effective (K, H) to (G, P) giving the geometry at u_0 with $\Sigma_J^{\mathfrak{k}} \neq \emptyset$.

In particular, if $\Sigma_J^{\text{inf}(u_0)} \neq \emptyset$, then there is $u_0 \in \mathcal{G}$ such that

$$\Sigma_J^{\text{inf}(u_0)} = \{s_J \exp Z : Z \in (\mathfrak{a}_{u_0})_+\}.$$

Proof. We know from the Lemma 3.5 that there is u_0 such that if $\Sigma_J^{\mathfrak{k}} \neq \emptyset$, then $s_J \in \Sigma_J^{\mathfrak{k}}$ holds. Since $s_J^{-1} \cdot \Sigma_J^{\mathfrak{k}} = \Sigma_{\text{id}_{\mathfrak{g}}^{-1}/\mathfrak{p}}^{\mathfrak{k}}$, the first claim follows from the Lemma 3.6. In particular, the second claim follows for the case $\mathfrak{k} = \text{inf}(u_0)$. \square

4. HOMOGENEOUS PARABOLIC GEOMETRIES AND GENERALIZED SYMMETRIES

Characterization of generalized symmetries. Now, we are ready to start the investigation of the generalized symmetries of a particular type $\Sigma(s_J)$ for $s_J \in Z(G_0)$. In particular, our results about automorphisms of type Σ_J provide several equivalent characterizations of homogeneous parabolic geometries admitting s_J -symmetries.

Theorem 4.1. *Let $(\pi : \mathcal{G} \rightarrow M, \omega)$ be a homogeneous parabolic geometry of type (G, P) and $s_J \in Z(G_0)$. Then there is $u_0 \in \mathcal{G}$ such that the following claims are equivalent:*

- (1) $\Sigma_J^{\text{inf}(u_0)} \neq \emptyset$, i.e., the geometry admits automorphisms of type Σ_J ,
- (2) $s_J \in A_{u_0}$, i.e., the geometry admits s_J -symmetries,
- (3) $\text{inf}(u_0)$ splits into eigenspaces of Ad_{s_J} in \mathfrak{g} and the components of the harmonic curvature such that $\lambda_J(\mu) \neq 0 \pmod{2\pi i}$ for the corresponding weight μ vanish at u_0 .

Proof. The description of $\Sigma_J^{\text{inf}(u_0)}$ at $u_0 \in \mathcal{G}$ in the Theorem 3.7 shows that (1) and (2) are equivalent. Further, claims (1) and (2) of the Lemma 3.4 show for the choice $\mathfrak{k} = \text{inf}(u_0)$ that (2) is equivalent with (3). \square

We proved in the Theorem 3.7 that the automorphisms of type Σ_J are in bijection with elements of $(\mathfrak{a}_{u_0})_+$. The following Corollary of the Lemma 3.3 shows which elements of $(\mathfrak{a}_{u_0})_+$ correspond to generalized symmetries of type $\Sigma(s_J)$.

Corollary 4.2. *Let $(\pi : \mathcal{G} \rightarrow M, \omega)$ be a homogeneous parabolic geometry of type (G, P) such that $s_J \in A_{u_0}$. Then s_J -symmetries are in bijection with the elements of the set*

$$(\mathfrak{a}_{u_0})_+ \cap \{C(-\text{Ad}_{s_J}^{-1}(Y), Y) : Y \in \mathfrak{p}_+\},$$

where the operator C is given by the BCH-formula.

Generalized symmetries and harmonic curvature. Let us now describe how the remaining curvature restricts $\Sigma_J^{\text{inf}(u_0)}$ and vice versa. There is the following action of \mathfrak{a}_{u_0} on the harmonic curvature for each one-parameter group of automorphisms ϕ_t generated by $X \in \mathfrak{a}_{u_0}$:

$$0 = \frac{d}{dt} \Big|_{t=0} \kappa_H(u_0) = \frac{d}{dt} \Big|_{t=0} \kappa_H(\phi_t(u_0)) = X \cdot \kappa_H(u_0).$$

This action together with the results in [10] allows us to formulate the following Theorem describing the relations between $(\mathfrak{a}_{u_0})_+$ and $\kappa_H(u_0)$.

Theorem 4.3. *Let $(\pi : \mathcal{G} \rightarrow M, \omega)$ be a homogeneous parabolic geometry of type (G, P) and assume $(\mathfrak{a}_{u_0})_+ \neq \emptyset$. Then the following facts hold:*

- (1) Denote

$$\mathfrak{a}_0(\kappa_H(u_0)) := \{X \in \mathfrak{g}_0 : X \cdot \kappa_H(u_0) = 0\},$$

$$\mathfrak{a}_i(\kappa_H(u_0)) := \{X \in \mathfrak{g}_i : [X, \mathfrak{g}_{-1}] \subset \mathfrak{a}_{i-1}(\kappa_H(u_0))\}.$$

Then $\text{gr}(\mathfrak{a}_{u_0}/(\mathfrak{a}_{u_0})_+) \subset \mathfrak{a}_0(\kappa_H(u_0))$ and $\text{gr}(\mathfrak{a}_{u_0}^i/\mathfrak{a}_{u_0}^{i+1}) \subset \mathfrak{a}_i(\kappa_H(u_0))$, where gr is the chosen identification of $\mathfrak{g}^i/\mathfrak{g}^{i+1}$ with \mathfrak{g}_i .

- (2) For an arbitrary component μ of the harmonic curvature in $H^2(\mathfrak{g}_-, \mathfrak{g})$ with non-trivial intersection with $\kappa_H(u_0)$ holds:

$$\dim(\mathrm{gr}(\mathfrak{a}_{u_0}^i/\mathfrak{a}_{u_0}^{i+1})) \leq \dim(\mathfrak{a}_i(\kappa_H(u_0))) \leq \dim(\mathfrak{a}_i(X^{-\mu})).$$

- (3) There is a simple restricted root γ of a positive height such that $\langle \gamma, \tilde{\mu}_i \rangle = 0$ for $\tilde{\mu}_i$ in $H^2(\mathfrak{p}_+, \mathfrak{g})$ corresponding to non-trivial components of $\kappa_H(u_0)$.
- (4) Assume $s_J \in A_{u_0}$. Then the eigenspaces of Ad_{s_J} in $(\mathfrak{a}_{u_0})_+$ can have eigenvalues only of the form $e^{\lambda_J(\alpha)}$ for restricted roots α spanned by roots γ satisfying the claim (3).

Proof. The first step to prove all the claims is to complexify \mathfrak{g} , \mathfrak{p} , κ_H , μ , G_0 and $(\mathfrak{a}_{u_0})_+$.

Since $(\mathfrak{a}_{u_0})_+$ acts trivially on the harmonic curvature, we get $\mathrm{gr}(\mathfrak{a}_{u_0}/(\mathfrak{a}_{u_0})_+) \subset \mathfrak{a}_0(\kappa_H(u_0))$. Since $\mathfrak{g}^{-1}/\mathfrak{p} \subset \mathfrak{inf}(u_0)/\mathfrak{a}_{u_0}$, we get $\mathrm{gr}(\mathfrak{a}_{u_0}^i/\mathfrak{a}_{u_0}^{i+1}) \subset \mathfrak{a}_i(\kappa_H(u_0))$ for all i and we proved the claim (1).

Then we can apply the results from [10]. Namely, the Proposition 3.1.1 implies the claim (2), and the Theorem 3.3.3 implies the claim (3).

We will use the claim (2) of the Lemma 3.4 to modify the Proof of [10, Proposition 3.1.1] in order to prove the claim (4). We again consider the complexified situation and let Z belong to $\mathrm{gr}(\mathfrak{a}_{u_0}^i/\mathfrak{a}_{u_0}^{i+1}) \subset \mathfrak{a}_i(\kappa_H(u_0))$. Then the components of Z in different eigenspaces of Ad_{s_J} are in $\mathfrak{a}_i(\kappa_H(u_0))$, too. Thus we only have to prove that the eigenspaces of Ad_{s_J} in $\mathfrak{a}_i(\kappa_H(u_0))$ satisfy the claim.

According to the proof of [10, Proposition 3.1.1], for any weight μ' in $H^2(\mathfrak{g}_-, \mathfrak{g})$ in a non-trivial component μ of $\kappa_H(u_0)$, there is a sequence g_i of elements of complexified G_0 such that $g_i \cdot \mu'$ converges to $-\mu$. Thus $g_i \cdot \mathfrak{a}_i(\kappa_H(u_0))$ converges to a subset of $\mathfrak{a}_i(X^{-\mu})$, which is described by [10, Theorem 3.3.3]. The consequence of the description of $\mathfrak{a}_i(X^{-\mu})$ is that Ad_{s_J} has only the claimed eigenvalues on $\mathfrak{a}_i(X^{-\mu})$.

We modify the proof of [10, Proposition 3.1.1] to prove that the dimension of the eigenspace of Ad_{s_J} in $\mathfrak{a}_i(\kappa_H(u_0))$ is lower than the dimension of that eigenspace in $\mathfrak{a}_i(X^{-\mu})$. Precisely, the consequence of claim (2) of the Lemma 3.4 is that we can discuss each eigenspace of Ad_{s_J} separately, and the argumentation in the proof [10, Proposition 3.1.1] is still valid. Thus the dimension of the eigenspace of Ad_{s_J} in $\mathfrak{a}_i(g_i \cdot X^\mu)$ does not drop, and the claim (4) follows. \square

Remark 1. There are tables presented in [10] giving the classification of possible roots γ for components of the harmonic curvature of simple complex parabolic geometries. We will complete the list for the real cases of our interest in the last section.

The above Theorem has a simple but important consequence.

Corollary 4.4. *Let $(\pi : \mathcal{G} \rightarrow M, \omega)$ be a non-flat homogeneous parabolic geometry of type (G, P) , and assume that $\lambda_J(\gamma) = 0 \pmod{2\pi i}$ for all simple restricted roots satisfying the claim (3) of the Theorem 4.3. Then there is at most one s_J -symmetry at any point.*

5. IMPORTANT CLASSES OF GENERALIZED SYMMETRIES OF HOMOGENEOUS GEOMETRIES

Automorphisms with higher order fixed points. The first natural action on $T_x^{-1}M$ we can consider is simply the identity. In such case, the Theorem 3.7 states that automorphisms of type Σ_{id} on a homogeneous parabolic geometry are precisely the automorphisms with higher order fixed points investigated in [3]. The existence of s_{id} -symmetry provides no new informations about the parabolic geometry, because the only s_{id} -symmetry is the identity on M . Thus the harmonic

curvature can only be restricted by the number of automorphisms of type Σ_{id} as in Theorem 4.3.

Symmetries of parabolic geometries. The second most natural action on $T_x^{-1}M$ we can consider is $-\text{id}$. We already investigated geometries with automorphisms of type $\Sigma_{-\text{id}}$ in [4, 6, 15, 16] and called them symmetries in these articles. In this article, we will call them *usual symmetries* to distinguish them from the other types of generalized symmetries. Moreover, for comparison with the other types of generalized symmetries, it is convenient to denote J as tuple $(-, \dots, -)$ representing the action $-\text{id}$ of J on the root spaces of each simple restricted root in Ξ . It is simple to decide about the existence of J for a given parabolic geometry according to the Proposition 3.2.

Proposition 5.1. *Let \mathfrak{g} be a semisimple Lie algebra, and let \mathfrak{p} be a parabolic subalgebra of \mathfrak{g} corresponding to $\Xi = \{i_1, \dots, i_j\}$, which does not contain any simple ideal of \mathfrak{g} . Then there exists a type (G, P) with $s_{(-, \dots, -)} \in Z(G_0)$.*

We can describe the eigenspaces of $\text{Ad}_{s_{(-, \dots, -)}}$ in a fairly simple way as follows.

Proposition 5.2. *Let $(\mathcal{G} \rightarrow M, \omega)$ be a parabolic geometry of type (G, P) such that $s_{(-, \dots, -)} \in Z(G_0)$. Then the following facts hold:*

- (1) *The -1 -eigenspace of Ad_{s_J} in \mathfrak{g} equals to $\sum_{i \text{ odd}} \mathfrak{g}_i$, and the 1 -eigenspace of Ad_{s_J} in \mathfrak{g} equals to $\sum_{i \text{ even}} \mathfrak{g}_i$.*
- (2)

$$\mathfrak{p}_{\text{fix}}^{s_{(-, \dots, -)}} = \sum_{i > 0 \text{ even}} \mathfrak{g}_i.$$

- (3) *A usual symmetry is $s_{(-, \dots, -)}$ -symmetry if and only if the symmetry is involutive. In particular,*

$$\Sigma(s_{(-, \dots, -)}) = \{s_{(-, \dots, -)} \exp(Z) : Z \in \sum_{i > 0 \text{ odd}} \mathfrak{g}_i\}.$$

Proof. The first claim follows from a simple computation with $\lambda_{(-, \dots, -)}$, and then the second claim follows, too.

If $s_{(-, \dots, -)} \exp(Z) s_{(-, \dots, -)} \exp(Z) = \text{id}$ for $Z \in \mathfrak{p}_+$, then $\text{Ad}_{s_{(-, \dots, -)}}(Z) = -Z$, i.e., $Z \in \sum_{i > 0 \text{ odd}} \mathfrak{g}_i$. Thus

$$s_{(-, \dots, -)} \exp(Z) = \exp\left(-\frac{1}{2}Z\right) s_{(-, \dots, -)} \exp\left(\frac{1}{2}Z\right) \in \Sigma(s_{(-, \dots, -)})$$

is $s_{(-, \dots, -)}$ -symmetry. Conversely, it is clear that $p^{-1} s_{(-, \dots, -)} p \in \Sigma(s_{(-, \dots, -)})$ is involutive. In particular, the description of $\Sigma(s_{(-, \dots, -)})$ follows. \square

Thus the Theorem 4.1 can be rephrased due to the previous results as follows.

Theorem 5.3. *Let $(\mathcal{G} \rightarrow M, \omega)$ be a homogeneous parabolic geometry of type (G, P) such that $s_J \in Z(G_0)$ for $J = (-, \dots, -)$. Then there is $u_0 \in \mathcal{G}$ such that the following facts are equivalent:*

- (1) *There is a usual symmetry at one (and thus at each) point of M .*
- (2) *There is an involutive usual symmetry at one (and thus at each) point of M .*
- (3) *The space $\text{inf}(u_0)$ splits into $\sum_{i \text{ odd}} \mathfrak{g}_i$ and $\sum_{i \text{ even}} \mathfrak{g}_i$, and the non-zero components of the harmonic curvature have even homogeneity.*

In particular, there is the following consequence of the condition on the harmonic curvature.

Corollary 5.4. *If a homogeneous parabolic geometry does not admit a harmonic curvature of even homogeneity, and there is a usual symmetry at one point of M , then the whole curvature vanishes.*

The classification of parabolic geometries with \mathfrak{g} simple that admit a harmonic curvature of even homogeneity can be found in the tables in the last Section.

Let us point out some important consequences of the classification in the tables in the last Section.

Proposition 5.5. *The following facts hold for homogeneous parabolic geometries:*

- (1) *There is more than one usual symmetry if and only if there is more than one involutive usual symmetry.*
- (2) *If there is no simple restricted root γ of positive height such that $\langle \gamma, \tilde{\mu}_i \rangle = 0$ for $\tilde{\mu}_i$ in $H^2(\mathfrak{p}_+, \mathfrak{g})$ corresponding to non-trivial components of $\kappa_H(u_0)$, then the non-flat homogeneous parabolic geometry has at most one (involutive) usual symmetry at each point.*
- (3) *If $|k| \leq 2$ and \mathfrak{g} is simple, then the non-flat homogeneous parabolic geometry has at most one (involutive) usual symmetry at each point.*

Proof. It follows from the Proposition 5.2 that there is only one involutive usual symmetry if and only if $(\mathfrak{a}_{u_0})_+ \subset \mathfrak{p}_{fix}^{s(-, \dots, -)} = \sum_{i>0 \text{ even}} \mathfrak{g}_i$, but this is not possible for obvious reasons. Thus the claim (1) follows.

The claim (2) follows from the Theorem 4.3. The claim (3) then follows from the tables in the last Section. \square

In particular, this Proposition extends the known restrictions on the number of usual symmetries to all AHS-structures and parabolic contact structures.

The homogeneous parabolic geometries with a unique usual symmetry at each point can be immediately related to the classical symmetric and reflexion spaces, see [11].

Theorem 5.6. *Let $(\mathcal{G} \rightarrow M, \omega)$ be a homogeneous parabolic geometry with a single (involutive) usual symmetry at one (and thus at each) point of M . Then M with these symmetries has a structure of a homogeneous reflexion space, i.e., it is a correspondence space to a symmetric space.*

Proof. We know that $ps_{(-, \dots, -)}p^{-1}$ is an automorphisms of type $\Sigma_{(-, \dots, -)}$ for any $p \in A_{u_0}$. Then it is clear that $s_{(-, \dots, -)} \in Z(A_{u_0})$ due to the uniqueness of the symmetry and thus the result follows from [5, Theorem 3.4.4]. \square

Remark 2. We recall that although the group generated by usual symmetries acts transitively on M , the whole $\text{Aut}(\mathcal{G}, \omega)$ does not have to act effectively on the symmetric space from the previous Theorem.

Let us briefly discuss the semisimple case. For arbitrary semisimple Lie algebra \mathfrak{g} , we will write $\mathfrak{g} = \sum_{l=1}^r \mathfrak{g}^{(l)}$, where $\mathfrak{g}^{(l)}$ are simple factors of \mathfrak{g} , which are naturally $|k_l|$ -graded. Then $\mu^{\mathfrak{g}} = \bigoplus_{l=1}^r \mu^{\mathfrak{g}^{(l)}}$, i.e., the component of the harmonic curvature (i, j) decomposes into factors $(i, j)^l$. Thus the components of the following types can appear, where $1 \leq l_1, l_2, l_3 \leq j$ are different, $b = 2 \frac{\langle \alpha_j, \alpha_i \rangle}{\langle \alpha_i, \alpha_i \rangle}$ and c is the homogeneity of $X^{\alpha_i} \otimes X^{s_{\alpha_i}(-\mu^{\mathfrak{g}^{(l_3)}})} \in \mathfrak{p}_+ \otimes \mathfrak{g}$:

TABLE 1. Possible types of components of the harmonic curvature for the semisimple Lie algebra \mathfrak{g}

	$(i, j)^l$	homogeneity	restrictions
(1)	$\mathfrak{g}_{\alpha_i} \subset \mathfrak{g}^{(l_1)}, \mathfrak{g}_{\alpha_j} \subset \mathfrak{g}^{(l_2)}, l = l_3$	1	$k_{l_3} = 1$
(2)	$\mathfrak{g}_{\alpha_i} \subset \mathfrak{g}^{(l_1)}, \mathfrak{g}_{\alpha_j} \subset \mathfrak{g}^{(l_1)}, l = l_3$	$1 - b - k_{l_3}$	$j \notin \Xi _{\mathfrak{g}^{(l_1)}}, b \neq 0, k_{l_3} < 1 - b$
(3)	$\mathfrak{g}_{\alpha_i} \subset \mathfrak{g}^{(l_1)}, \mathfrak{g}_{\alpha_j} \subset \mathfrak{g}^{(l_1)}, l = l_3$	$2 - b - k_{l_3}$	$j \in \Xi _{\mathfrak{g}^{(l_1)}}, k_{l_3} < 2 - b$
(4)	$\mathfrak{g}_{\alpha_i} \subset \mathfrak{g}^{(l_1)}, \mathfrak{g}_{\alpha_j} \subset \mathfrak{g}^{(l_2)}, l = l_1$	$1 + c$	$-1 < c$
(5)	$\mathfrak{g}_{\alpha_i} \subset \mathfrak{g}^{(l_1)}, \mathfrak{g}_{\alpha_j} \subset \mathfrak{g}^{(l_1)}, l = l_1$		

There is the following trivial but crucial observation about usual symmetries in the semisimple case.

Lemma 5.7. *If \mathfrak{g} is semisimple, then there generically is more than one (involutive) usual symmetry at a single point of M . It suffices to take a simple factor $\mathfrak{g}^{(l)}$ which does not contribute to the κ_H according to the Table 1.*

So we will discuss only the usual symmetries, which are different after restricting to simple factors $\mathfrak{g}^{(l_i)}$ which contribute to κ_H . Let us present here facts about possible components of the harmonic curvature of homogeneous parabolic geometries with $\Sigma_{(-, \dots, -)}^{\text{inf}(u_0)} \neq \emptyset$ according to the line in the Table 1.

- (1) The component always vanishes due to its homogeneity.
- (2) The component vanishes with the exception of the cases given in the following table (and their complexifications).

$\mathfrak{g}^{(l_1)}$	$\Xi _{\mathfrak{g}^{(l_1)}}$ contains	restrictions
$\mathfrak{g}_2(2)$	$\{1\}$	$k_{l_3} = 2$
$\mathfrak{f}_4(4)$	$\{2\}$	$k_{l_3} = 1$
$\mathfrak{so}(n, n+1)$	$\{n\}$	$k_{l_3} = 1$
$\mathfrak{sp}(2n, \mathbb{R})$	$\{n-1\}$	$k_{l_3} = 1$

In particular, if $\Xi|_{\mathfrak{g}^{(l_3)}}$ contains ℓ such that $\langle \mu^{\mathfrak{g}^{(l_3)}}, \alpha_\ell \rangle = 0$, then non-flat homogeneous parabolic geometries of the corresponding type admit more than one usual symmetry at each point.

- (3) The component vanishes with the exception of the cases given in the following table (and their complexifications).

$\mathfrak{g}^{(l_1)}$	$\Xi _{\mathfrak{g}^{(l_1)}}$ contains	restrictions
$\mathfrak{g}_2(2)$	$\{1, 2\}$	$k_{l_3} = 1, 3$
$\mathfrak{f}_4(4)$	$\{2, 3\}$	$k_{l_3} = 2$
$\mathfrak{so}(n, n+1)$	$\{n-1, n\}$	$k_{l_3} = 2$
$\mathfrak{sp}(2n, \mathbb{R})$	$\{n-1, n\}$	$k_{l_3} = 2$
arbitrary	$\{i, j\}$	$b = 1, k_{l_3} = 1$

In particular, if $\Xi|_{\mathfrak{g}^{(l_3)}}$ contains ℓ such that $\langle \mu^{\mathfrak{g}^{(l_3)}}, \alpha_\ell \rangle = 0$, then non-flat homogeneous parabolic geometries of the corresponding type admit more than one usual symmetry at each point.

- (4) The component vanishes with the exception of the cases satisfying $c = 1$, which are exactly projective and contact projective types. Then, if there is $\ell \in \Xi|_{\mathfrak{g}^{(l_2)}}$ such that $\sigma^*(\alpha_\ell) \neq \pm \alpha_j$ and $\langle \alpha_\ell, \alpha_j \rangle = 0$, then non-flat homogeneous parabolic geometries of the corresponding type admit more than one usual symmetry at each point.

(5) The component is an element of $H^2(\mathfrak{g}_-^{(l_1)}, \mu^{\mathfrak{g}^{(l_1)}})$, and results from the simple case apply.

Generalized symmetries of order 2 and parabolic geometries with weak para-complex structures. Let us now consider automorphisms of type Σ_J , where J is a natural weak para-complex structure on $T_x^{-1}M$, i.e., $J^2 = \text{id}$ holds. We will show that there can be many different para-complex structures on $T_x^{-1}M$, and it is convenient to denote J as tuple with entries $+$ or $-$ that indicate, whether the action of J is id or $-\text{id}$ on root spaces of each simple restricted root in Ξ . The possible para-complex structures J for a given parabolic geometry are determined by the Proposition 3.2.

Proposition 5.8. *Let \mathfrak{g} be a semisimple Lie algebra, and let \mathfrak{p} be a parabolic subalgebra of \mathfrak{g} corresponding to $\Xi = \{i_1, \dots, i_j\}$, which does not contain any simple ideal of \mathfrak{g} . Then there exists a type (G, P) with $s_J \in Z(G_0)$ such that the corresponding J is a weak almost para-complex structure on $T^{-1}M$ if and only if $j > 2$, or $j = 2$ and $\sigma^*(\alpha_{i_1}) \neq \alpha_{i_2}$.*

All possible para-complex structures J correspond to decompositions of Ξ into $\Xi^- = \{i_\ell \in \Xi : J_\ell = -\}$ and $\Xi^+ = \{i_\ell \in \Xi : J_\ell = +\}$, which are stable under the complex conjugation σ^ .*

The structure of eigenspaces of Ad_{s_J} is more complicated in this case. We have to consider the bi-grading $\mathfrak{g}_{(a,b)} \subset \mathfrak{g}_{a+b}$ of \mathfrak{g} with respect to Ξ^- and Ξ^+ . Then we can characterize the P -conjugacy class $\Sigma(s_J)$ as follows.

Proposition 5.9. *Let $(\mathcal{G} \rightarrow M, \omega)$ be a parabolic geometry of type (G, P) such that $s_J \in Z(G_0)$ for the para-complex structure J . Then the following facts hold:*

- (1) *The -1 -eigenspace of Ad_{s_J} in \mathfrak{g} is $\sum_{a \text{ odd}} \mathfrak{g}_{(a,b)}$, and the 1 -eigenspace of Ad_{s_J} in \mathfrak{g} is $\sum_{a \text{ even}} \mathfrak{g}_{(a,b)}$.*
- (2)

$$\mathfrak{p}_{fix}^{s_J} = \sum_{a+b>0, a \text{ even}} \mathfrak{g}_{(a,b)}.$$

- (3) *An automorphism of type Σ_J is s_J -symmetry if and only if the symmetry is involutive. In particular,*

$$\Sigma(s_J) = \{s_J \exp(Z) : Z \in \sum_{a+b>0, a \text{ odd}} \mathfrak{g}_{(a,b)}\}.$$

Proof. The first claim follows from a simple computation with λ_J on $\mathfrak{g}_{(a,b)}$, and then the second claim is clear.

If $s_J \exp(Z) s_J \exp(Z) = \text{id}$ for $Z \in \mathfrak{p}_+$, then $\text{Ad}_{s_J}(Z) = -Z$ holds, i.e., $Z \in \sum_{a+b>0, a \text{ odd}} \mathfrak{g}_{(a,b)}$. Thus

$$s_J \exp(Z) = \exp(-\frac{1}{2}Z) s_J \exp(\frac{1}{2}Z) \in \Sigma(s_J)$$

is s_J -symmetry. Conversely, it is clear that $p^{-1} s_J p \in \Sigma(s_J)$ is involutive. In particular, the description of $\Sigma(s_J)$ follows. \square

The condition (3) of Theorem 4.1 implies the following.

Proposition 5.10. *Let $(\mathcal{G} \rightarrow M, \omega)$ be a homogeneous parabolic geometry of type (G, P) such that $s_J \in Z(G_0)$ for the para-complex structure J . Then the bi-homogeneity (a, b) of each component κ_H has a even.*

The classification of parabolic geometries with \mathfrak{g} simple that admit non-trivial harmonic curvatures can be found in the tables in the last Section.

The statement analogous to the Proposition 5.5 is more complicated, because we have to distinguish between the number of automorphisms of type Σ_J and the number of s_J -symmetries.

Proposition 5.11. *The following facts hold for non-flat homogeneous parabolic geometries:*

- (1) *If there is no simple restricted root γ of positive height such that $\langle \gamma, \tilde{\mu}_i \rangle = 0$ for $\tilde{\mu}_i$ in $H^2(\mathfrak{p}_+, \mathfrak{g})$ corresponding to non-trivial components of $\kappa_H(u_0)$, then there is at most one automorphism of type Σ_J at each point.*
- (2) *If there is no simple restricted root γ such that $\mathfrak{g}_\gamma \in \mathfrak{g}_{(1,0)}$ and $\langle \gamma, \tilde{\mu}_i \rangle = 0$ for $\tilde{\mu}_i$ in $H^2(\mathfrak{p}_+, \mathfrak{g})$ corresponding to non-trivial components of $\kappa_H(u_0)$, then there is at most one s_J -symmetry at each point.*

Proof. The claim (1) follows from the Theorem 4.3, and the claim (2) is a simple consequence of the Corollary 4.4. \square

Let us point out the important consequence of the Theorem 4.3.

Theorem 5.12. *Let $(\mathcal{G} \rightarrow M, \omega)$ be a homogeneous parabolic geometry with a single s_J -symmetry at one (and thus at each) point of M . Then M with these s_J -symmetries has a structure of a homogeneous reflexion space, i.e., it is a correspondence space to a non-effective homogeneous symmetric space.*

Proof. We know that $ps_{(-, \dots, -)}p^{-1}$ is an automorphisms of type $\Sigma_{(-, \dots, -)}$ for any $p \in A_{u_0}$. Then it is clear that $s_{(-, \dots, -)} \in Z(A_{u_0})$ due to the uniqueness of the s_J -symmetry and thus the result follows from [5, Theorem 3.4.4]. However, since the group generated by s_J -symmetries does not have to act transitively on M , then $\text{Aut}(\mathcal{G}, \omega)$ does not have to act effectively on the symmetric space. \square

Since the 1-eigenspace of Ad_{s_J} is too big, there are too many possible cases we have to distinguish, when \mathfrak{g} is semisimple. So although it is possible to derive the analogous restrictions as in the case of usual symmetries, we will not discuss them here explicitly. We only recall that the Lemma 5.7 is still valid for this type of generalized symmetries.

Generalized symmetries of order 4 and parabolic geometries with complex structures. Finally, we consider automorphisms of type Σ_J , where J is a natural complex structure on $T_x^{-1}M$, i.e., $J^2 = -\text{id}$ holds. We will show that there can be many different complex structures on $T_x^{-1}M$, and it is convenient to denote J as tuple with entries i or $-i$ that indicate, whether the action of J is $i \cdot \text{id}$ or $-i \cdot \text{id}$ on the root space of each simple restricted root in Ξ . The possible complex structures J for a given parabolic geometry are determined by the Proposition 3.2 as follows.

Proposition 5.13. *Let \mathfrak{g} be a semisimple Lie algebra, and let \mathfrak{p} be a parabolic subalgebra of \mathfrak{g} corresponding to $\Xi = \{i_1, \dots, i_j\}$, which does not contain any simple ideal of \mathfrak{g} . Then there exists a type (G, P) with $s_J \in Z(G_0)$ such that J is an almost complex structure on $T^{-1}M$ if and only if $\sigma^*(\alpha_{i_l}) \neq -\alpha_{i_l}$ for $l = 1, \dots, j$.*

All possible complex structures J correspond to decompositions of Ξ into $\Xi^- = \{i_l \in \Xi : J_l = -i\}$ and $\Xi^+ = \{i_l \in \Xi : J_l = i\}$ such that $\sigma^(\Xi^-) = \Xi^+$.*

These generalized symmetries can be naturally related to the previous two classes of generalized symmetries.

Corollary 5.14. *Let $(\mathcal{G} \rightarrow M, \omega)$ be a homogeneous parabolic geometry with s_J -symmetry at one (and thus at each) point of M . Then there is $s_J^2 = s_{(-, \dots, -)}$ -symmetry at one (and thus at each) point of M . In general, the composition $J \circ J'$ of two complex structures J and J' is a weak para-complex structure.*

In other words, if there is at least one complex structure, then we can use the results from the previous two cases to get all of them.

There again is the bi-grading $\mathfrak{g}_{(a,b)} \subset \mathfrak{g}_{a+b}$ of \mathfrak{g} with respect to Ξ^- and Ξ^+ . The eigenspaces of Ad_{s_J} can be described as follows.

Proposition 5.15. *Let $(\mathcal{G} \rightarrow M, \omega)$ be a parabolic geometry of type (G, P) such that $s_J \in Z(G_0)$ for the complex structure J . Then the following facts hold:*

- (1) *The i^c -eigenspace of Ad_{s_J} in \mathfrak{g} is $\sum_{3a+b=c \pmod 4} \mathfrak{g}_{(a,b)}$. In particular, the 1-eigenspace of Ad_{s_J} in \mathfrak{g} is $\sum_{3a+b=0 \pmod 4} \mathfrak{g}_{(a,b)}$.*
- (2)

$$\mathfrak{p}_{fix}^{s_J} = \sum_{a+b>0, 3a+b=0 \pmod 4} \mathfrak{g}_{(a,b)}.$$

Proof. The first claim follows from a simple computation with λ_J on $\mathfrak{g}_{(a,b)}$, and then the second claim is clear. \square

The condition (3) of the Theorem 4.1 implies the following.

Proposition 5.16. *Let $(\mathcal{G} \rightarrow M, \omega)$ be a homogeneous parabolic geometry of type (G, P) such that $s_J \in Z(G_0)$ for the complex structure J . Then the bi-homogeneity (a, b) of each component of κ_H satisfies $3a + b = 0 \pmod 4$.*

The classification of parabolic geometries with \mathfrak{g} simple that admit such harmonic curvatures can be found in the tables in the last Section.

Proof. Since $(-i)^a(i)^b = 1$ if and only if $3a + b = 0 \pmod 4$, the first claim is clear.

In order to get the results in the tables, we discuss the condition on the bi-homogeneity in more detail. Since the homogeneity has to be even, the possible homogeneities are 2 with $(a, b) = (3, -1), (1, 1), (-1, 3)$, and 4, which appear only in special dimensions and can be computed directly.

If \mathfrak{g} is simple, but the complexification $\mathfrak{g}_{\mathbb{C}}$ is not simple, then $\mathfrak{g}_{\mathbb{C}} = \mathfrak{g} \oplus \mathfrak{g}$ with the highest root $\mu^{\mathfrak{g}_{\mathbb{C}}} = \mu^{\mathfrak{g}} + (\mu^{\mathfrak{g}})'$, where we denote by α' the roots of the other copy of \mathfrak{g} . In particular, $\sigma^*(\alpha) = \alpha'$. So J can be chosen arbitrarily on each root $\alpha \in \Xi$, and the values on $\alpha' \in \Xi$ are determined by the above choice. Then, according to the line in the Table 1, we have the following facts for corresponding components of the harmonic curvature of homogeneous parabolic geometries with $s_J \in A_{u_0}$:

- (1) These components cannot appear.
- (2) These components cannot appear, because $b \leq k$.
- (3) These components cannot appear, because $b < k$.
- (4) These components have to have homogeneity 2, i.e., they arise only on complex projective and complex contact projective geometries.
- (5) There are two possibilities for the homogeneity of the component (i, j) . If $\alpha_j \notin \Xi$, then the homogeneity has to be 4. If $\alpha_j \in \Xi$, then the homogeneity can be 2 or 4, and we simply check the bi-homogeneity for each of them.

If $\mathfrak{g}_{\mathbb{C}}$ is simple, we have to distinguish if $\sigma^*(\alpha_i) = \alpha_j$ or not for the component of the harmonic curvature (i, j) . If not, then the possibilities are the same as in (5). Otherwise, the bi-homogeneity has to be $(1, 1)$ or $(2, 2)$, which is possible only in the $\mathfrak{g} = \mathfrak{su}(p, q)$ -case. \square

The Proposition 5.5 naturally holds in this case, too, and we get the following result in the same way as for usual symmetries.

Theorem 5.17. *Let $(\mathcal{G} \rightarrow M, \omega)$ be a homogeneous parabolic geometry with a single s_J -symmetry at one (and thus at each) point of M . Then M with these $(s_J)^2$ -symmetries has a structure of a homogeneous reflexion space, i.e., it is a correspondence space to a complex symmetric space. This is particularly true, when \mathfrak{g} is simple, $|k| \leq 2$ and the geometry is non-flat.*

Again, there are too many choices involved for the action of J , when \mathfrak{g} is semisimple. So although we can derive the analogous restrictions as in the case of usual symmetries, we will not discuss them here. We only recall that the Lemma 5.7 is still valid for this type of generalized symmetries.

6. TWO GENERAL CONSTRUCTIONS OF EXAMPLES

Extensions of correspondence spaces over symmetric spaces. Let us recall the construction of symmetric parabolic geometries from [4, 5], which particularly allows to construct symmetries of all types discussed in the Section 5. The construction can be summarized in the following steps:

- (1) We take the effective pair (K, H) such that there is $h \in H$ such that H is contained in the centralizer of h in K , and K/H is connected.
- (2) We find the extension (i, α) of (K, H) to (G, P) such that $i(h) \in \Sigma(s_J)$ for some $s_J \in Z(G_0)$.
- (3) Then $\mathcal{F}_\alpha(K \rightarrow K/H, \omega_K)$ will be s_J -symmetric, where $J = \underline{\text{Ad}}_{i(h)}|_{\mathfrak{g}^{-1/\mathfrak{p}}}$.

If $h^2 = \text{id}$, then $K/H \rightarrow K/Z_K(h)$ is a correspondence space over the symmetric space $K/Z_K(h)$, where $Z_K(h)$ is the centralizer of h in K , and the corresponding generalized symmetries cover the symmetries on $K/Z_K(h)$. In particular, one can start with the known classifications of symmetric spaces and homogeneous bundles over them.

However, without any distinguished properties of the elements $h \in H$, there is no classification of the homogeneous spaces K/H , and we cannot use the construction explicitly.

There are many examples of non-flat homogeneous parabolic geometries constructed by the construction from symmetric spaces in [4, 5]. We present one family of them here.

Example 2 (Extension $\alpha : \mathfrak{so}(3) \rightarrow \mathfrak{su}(2, 1)$). Consider the one-parameter class of linear maps $\alpha_t : \mathfrak{so}(3) \rightarrow \mathfrak{su}(2, 1)$ for $t \geq 1$ given by

$$\alpha_t(x_1, x_2, x_3) = \begin{pmatrix} \frac{1+t^4}{8t^2}ix_3 & * & \frac{-(15t^8-34t^4+15)}{128t^4}ix_3 \\ tx_1 + \frac{i}{t}x_2 & -\frac{1+t^4}{4t^2}ix_3 & \frac{-3t^4+5}{16t}x_1 + \frac{5t^4-3}{16t^3}ix_2 \\ 2ix_3 & * & \frac{1+t^4}{8t^2}ix_3 \end{pmatrix},$$

where the entries denoted by $*$ are determined by the structure of $\mathfrak{su}(2, 1)$, and (x_1, x_2, x_3) represents the element

$$\begin{pmatrix} 0 & x_3 & -x_1 \\ -x_3 & 0 & -x_2 \\ x_1 & x_2 & 0 \end{pmatrix} \in \mathfrak{so}(3).$$

The curvature of the homogeneous parabolic geometry given at u_0 by an extension with such α_t is of the form

$$\kappa((x_1, x_2, x_3), (y_1, y_2, y_3))(u_0) = \begin{pmatrix} 0 & * & 0 \\ 0 & 0 & \frac{3(1-t^8)}{16t^5}(y_3x_2 - x_3y_2) + \frac{3(1-t^8)}{16t^3}i(y_3x_1 - x_3y_1) \\ 0 & 0 & 0 \end{pmatrix},$$

where the entries denoted by $*$ are determined by the structure of $\mathfrak{su}(2, 1)$, and κ vanishes for $t = 1$.

Thus the map α_1 is a Lie algebra homomorphism and thus determines the Lie group homomorphism $\iota_1 : K := Spin(3) \rightarrow G := PSU(2, 1)$. Let H be the preimage of $\iota_1(Spin(3)) \cap P_{1,2}$. Then $(\iota_1|_H, \alpha_t)$ for arbitrary $t \geq 1$ is an extension of

($Spin(3), H$) to $(PSU(2, 1), P_{1,2})$ giving a regular normal parabolic geometry with the above curvature. We recall that $(PSU(2, 1), P_{1,2})$ is the type of CR-geometries.

It follows from the construction that

$$s_{(-,-)} = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$

is the usual symmetry for the geometry given by any of the above extensions.

Further, we can consider the element

$$s_{(i,-i)} := \begin{pmatrix} i & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & i \end{pmatrix} \in Z(G_0),$$

and it is easy to check that $\text{Ad}_{s_{(i,-i)}}\alpha_t(\mathfrak{so}(3)) \subset \alpha_t(\mathfrak{so}(3))$ and $\text{Ad}_{s_{(i,-i)}}\kappa(u_0) = \kappa(u_0)$ hold if and only if $t = 1$. Thus there is $s_{(i,-i)}$ -symmetry on the parabolic geometry of type $(PSU(2, 1), P_{1,2})$ given by the extension $(\iota_1|_H, \alpha_1)$, which is flat.

Deformations of nilpotent groups by the lowest weight of harmonic curvature. We introduce the construction, which generalizes the construction from [10] to the real case. The construction can be divided into the following steps:

- (1) We start with \mathfrak{g}_- of the chosen type (G, P) . (In general, we can start with the sum \mathfrak{b}_- of all negative root spaces.)
- (2) We choose a component of the harmonic curvature, which should not vanish. We know that the lowest weight vector $X^{-\mu}$ of the weight μ representing the component of the harmonic curvature can be seen as a map $X^{-\mu} : \mathfrak{g}_- \wedge \mathfrak{g}_- \rightarrow \mathfrak{g}$. In our examples, the map $X^{-\mu}$ has its image in \mathfrak{g}_- . (Generically, the image is in \mathfrak{b}_- , and there are several rank two geometries such that the image is in \mathfrak{p}_+ (for example see Example 2).)
- (3) We consider the Lie algebra \mathfrak{n} coinciding with \mathfrak{g}_- as a vector space together with the Lie bracket $[\cdot, \cdot]_{\mathfrak{g}_-} + X^{-\mu}$. The proof that \mathfrak{n} is a Lie algebra can be found in [10, Lemma 4.1.1]. The natural identification $\alpha_{\mathfrak{n}} : \mathfrak{n} \rightarrow \mathfrak{g}_-$ is an extension of $(\exp(\mathfrak{n}), \{e\})$ to (G, P) with the curvature $-X^{-\mu}$. (In general, $\mathfrak{b}_0 := \mathfrak{b}_- \cap \mathfrak{p}$ is nilpotent, and $\alpha_{\mathfrak{n}} : \mathfrak{n} \oplus \mathfrak{b}_0 \rightarrow \mathfrak{b}_-$ is an extension of $(\exp(\mathfrak{n} \oplus \mathfrak{b}_0), \exp(\mathfrak{b}_0))$ to (G, P) .)
- (4) We compute $(A_{u_0})_0$ following the Lemma 2.2, and we compute $(\mathfrak{a}(-X^{-\mu}))_+$ following the Theorem 4.3.
- (5) We describe $Z(G_0) \cap (A_{u_0})_0$ in order to find all types of generalized symmetries that the geometry admits. If $(\mathfrak{a}(-X^{-\mu}))_+ = 0$, then we end here. But if $(\mathfrak{a}(-X^{-\mu}))_+ \neq 0$, the corresponding infinitesimal automorphisms are not complete on $\exp(\mathfrak{n})$, and we need to modify our construction in the following way to get $(\mathfrak{a}_{u_0})_+ = (\mathfrak{a}(-X^{-\mu}))_+$:

We consider the decomposition of G/P into Schubert cells of the form

$$G/P = \bigsqcup_{w \in W^{\mathfrak{p}}} \exp(\mathfrak{b}_-) \tilde{w}.eP,$$

where $W^{\mathfrak{p}}$ is the Hasse graph of \mathfrak{p} in \mathfrak{g} , and \tilde{w} are the corresponding elements of G . Then we define the manifold M as

$$M := \bigsqcup_{w \in W^{\bar{\mathfrak{p}}}} \exp(\mathfrak{b}_-) \tilde{w}.eP,$$

where $W^{\bar{\mathfrak{p}}}$ is Hasse graph of $\bar{\mathfrak{p}}$ in $\bar{\mathfrak{g}} \subset \mathfrak{g}$, which consists of connected components of Dynkin diagram containing only the simple restricted roots α_i such that $\langle \alpha_i, \bar{\mu} \rangle = 0$ for the corresponding $\bar{\mu}$ in $H^2(\mathfrak{p}_+, \mathfrak{g})$ and $\bar{\mathfrak{p}} = \bar{\mathfrak{g}} \cap \mathfrak{p}$. The

following Proposition shows that there is a non-flat homogeneous parabolic geometry on M with the stabilizer $(A_{u_0})_0 \exp((\mathfrak{a}(-X^{-\mu}))_+)$.

Proposition 6.1. *There is a Lie group K with the Lie algebra $\mathfrak{n} \oplus \mathfrak{a}$ acting transitively on M , which is homotopically equivalent to a subgroup of G with the Lie algebra $\mathfrak{g}_- \oplus \mathfrak{a}$. Moreover, this determines an extension of (K, H) to (G, P) , where $H = (A_{u_0})_0 \rtimes \exp((\mathfrak{a}(-X^{-\mu}))_+)$ is the stabilizer of a point in M . The parabolic geometry over $M = K/H$ of type (G, P) given at u_0 by this extension is regular and normal, and K coincides with the automorphism group of the geometry.*

Proof. It follows from the results in [10, Section 3.3] that $(\mathfrak{a}(-X^{-\mu}))_+ = \bar{\mathfrak{p}}_+$, and thus, there is a subgroup K' of G with the Lie algebra $\mathfrak{g}_- \oplus \mathfrak{a}(-X^{-\mu})$ acting transitively on M . If we look on the intersection of K' with the Iwasawa decomposition of G , then it is clear that the group K' can be written as $\exp(\mathfrak{b}_-) \bar{A} \bar{C}$, where $N \bar{A} \bar{C}$ is the Iwasawa decomposition of the subgroup of K' with the Lie algebra $\bar{\mathfrak{g}}$, which is homotopically equivalent to K' .

Since $\exp(\mathfrak{b}_-)$ is simply connected, we can modify its Lie group structure to $\exp(\mathfrak{n} + \mathfrak{b}_0)$, and $K := \exp(\mathfrak{n} + \mathfrak{b}_0) \bar{A} \bar{C}$ is a Lie group with the Lie algebra $\mathfrak{n} \oplus \mathfrak{a}(-X^{-\mu})$ acting transitively on M , which is homotopically equivalent to K' .

So we get an extension of (K, H) to (G, P) , and since the only component of the curvature is harmonic, the geometry on K/H given by the extension is regular and normal. Since the action of any other element of P on the curvature is non-trivial, K coincides with the automorphism group of the geometry. \square

Now, we will present several examples of the above construction. We will discuss the details of the construction only in the first example. The remaining examples are analogous, and we will summarize here only the results.

Example 3 (Type $(\mathfrak{sl}(4, \mathbb{R}), \mathfrak{p}_{1,2,3})$ with curvature $(2, 1)$). (1) The pair $(\mathfrak{g}, \mathfrak{p}) = (\mathfrak{sl}(4, \mathbb{R}), \mathfrak{p}_{1,2,3})$ has \mathfrak{g}_- of the form

$$\mathfrak{g}_- = \mathfrak{b}_- = \begin{pmatrix} 0 & 0 & 0 & 0 \\ x_1 & 0 & 0 & 0 \\ x_2 & x_3 & 0 & 0 \\ x_5 & x_6 & x_4 & 0 \end{pmatrix}.$$

- (2) There are five possible components of the harmonic curvature, and we choose the component $(2, 1)$.
 (3) The curvature of the geometry given by the modified Lie bracket of \mathfrak{n} is of the form

$$\kappa((x_1, \dots, x_6), (x'_1, \dots, x'_6))(u_0) = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & x_3 x'_2 - x_2 x'_3 & 0 \end{pmatrix}.$$

- (4) We compute

$$(A_{u_0})_0 = \begin{pmatrix} x_7 & 0 & 0 & 0 \\ 0 & x_8 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & \frac{1}{x_7 x_8} \end{pmatrix},$$

$$(\mathfrak{a}_{u_0})_+ = \begin{pmatrix} 0 & x_9 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

- (5) It is obvious that $Z(G_0) \cap (A_{u_0})_0 = (A_{u_0})_0$, and \mathfrak{g}_{-1} decomposes into the eigenspaces with eigenvalues $\frac{x_8}{x_7}, \frac{1}{x_8}, \frac{1}{x_7x_8}$ of possible generalized symmetries. Thus there are generalized symmetries of all possible types according to the Table 5, namely:

$$s_{(-,-,-)} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix},$$

$$s_{(-,+,-)} = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix},$$

and $s_{(+,-,+)} = s_{(-,-,-)} \circ s_{(-,+,-)}$.

Since $(\mathfrak{a}_{u_0})_+ \neq 0$, there is infinitely many generalized symmetries of the first two types at each point, and there is infinitely many automorphisms of type $\Sigma_{(+,-,+)}$.

Remark 3. In this example, we get $\mathfrak{n} = \mathbb{R} \oplus \mathfrak{n}_{2,3,5}$, where \mathbb{R} is spanned by the x_1 -entry, and $\mathfrak{n}_{2,3,5}$ is the negative part of the grading of the famous parabolic geometry $(G_2(2), P_1)$ corresponding to the generic distribution of the growth rate $(2, 3, 5)$.

It is easy to check that $K = Gl(2, \mathbb{R}) \times \exp(\mathfrak{n}_{2,3,5}) \subset G_2(2)$, and M is homotopically equivalent to $S^1 \times \mathbb{R}^5$.

Example 4 (Type $(\mathfrak{so}(3, 5), \mathfrak{p}_{1,2})$ with curvature $(1, 2)$). In this example, the data for the construction are the following:

$$\mathfrak{g}_- = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ x_1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ x_2 & x_3 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -x_4 & -x_5 & 0 & -x_1 & -x_2 & -x_{10} & -x_8 \\ x_4 & 0 & -x_6 & 0 & 0 & -x_3 & -x_7 & -x_9 \\ x_5 & x_6 & 0 & 0 & 0 & 0 & 0 & 0 \\ x_{10} & x_7 & 0 & 0 & 0 & 0 & 0 & 0 \\ x_8 & x_9 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$\kappa((x_1, \dots, x_{10}), (x'_1, \dots, x'_{10}))(u_0) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & x_1x'_2 - x'_1x_2 & 0 & 0 & 0 & 0 & 0 \\ 0 & -x_1x'_2 + x'_1x_2 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$(A_{u_0})_0 = \begin{pmatrix} \pm x_{14}x_{15} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & x_{14} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & x_{15} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{\pm 1}{x_{14}x_{15}} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{x_{14}} & 0 & 0 & 0 \\ 0 & 0 & \frac{-x_{11}^2 - x_{12}^2}{2} & 0 & 0 & \frac{1}{x_{15}} & -x_{11} & -x_{12} \\ 0 & 0 & x_{11} & 0 & 0 & 0 & \cos(x_{13}) & -\sin(x_{13}) \\ 0 & 0 & x_{12} & 0 & 0 & 0 & \sin(x_{13}) & \cos(x_{13}) \end{pmatrix},$$

$$(\mathfrak{a}_{u_0})_+ = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & x_{16} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -x_{16} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

It is easy to check that $Z(G_0) \cap (A_{u_0})_0$ is given by $x_{11} = x_{12} = x_{13} = 0$ and $x_{15} = 1$, and \mathfrak{g}_{-1} decomposes into the eigenspaces with eigenvalues $\pm 1, x_{14}$ of possible generalized symmetries. Thus there are generalized symmetries of all possible types according to the Table 3, namely:

$$s_{(-,-)} := \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$s_{(+,-)} := \begin{pmatrix} -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

and $s_{(-,+)} := s_{(-,-)} \circ s_{(+,-)}$.

Since $(\mathfrak{a}_{u_0})_+ \neq 0$, there is infinitely many generalized symmetries of the first two types at any point, and there is infinitely many automorphisms of type $\Sigma_{(-,+)}$.

Remark 4. The Lie group K can be expressed in the following way: We decompose \mathfrak{n} as $\mathfrak{n} = \mathbb{R} \oplus \mathfrak{n}_{4,7,8,9}$, where \mathbb{R} is spanned by the x_3 -entry, and the part of A_{u_0} not contained in \bar{G} is the opposite parabolic subgroup $(P_1)^{op}$ in $PSO(1,3)$. Then we can write $K = PGL(2, \mathbb{R}) \rtimes (\exp(\mathfrak{n}_{4,7,8,9}) \times (P_1)^{op})$.

Example 5 (Type $(\mathfrak{sp}(8, \mathbb{R}), \mathfrak{p}_{1,2,4})$ with curvature $(2,1)$). In this example, the data for the construction are the following:

$$\mathfrak{g}_- = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ x_1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ x_6 & x_2 & 0 & 0 & 0 & 0 & 0 & 0 \\ x_7 & x_3 & 0 & 0 & 0 & 0 & 0 & 0 \\ x_{15} & x_{14} & x_{12} & x_{10} & 0 & -x_1 & -x_6 & -x_7 \\ x_{14} & x_{13} & x_{11} & x_8 & 0 & 0 & -x_2 & -x_3 \\ x_{12} & x_{11} & x_9 & x_4 & 0 & 0 & 0 & 0 \\ x_{10} & x_8 & x_4 & x_5 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$\kappa((x_1, \dots, x_{15}), (x'_1, \dots, x'_{15})) =$$

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & x_6 x'_2 - x_2 x'_6 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$(A_{u_0})_0 = \begin{pmatrix} x_{16} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{x_{17}^4}{x_{16}} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & x_{17} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & x_{19} & x_{18} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{x_{16}} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{x_{16}}{x_{17}^4} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{x_{17}} & \frac{-x_{19}}{x_{17} x_{18}} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{x_{18}} \end{pmatrix},$$

$$(\mathfrak{a}_{u_0})_+ = \begin{pmatrix} 0 & x_{20} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & x_{21} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -x_{20} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

It is easy to check that $Z(G_0) \cap (A_{u_0})_0$ is given by $x_{17} = x_{18}$ and $x_{19} = 0$, and \mathfrak{g}_{-1} decomposes into the eigenspaces with eigenvalues $\frac{x_{17}^4}{x_{16}^2}, \frac{x_{16}}{x_{17}^3}, \frac{1}{x_{17}^2}$ of possible generalized symmetries. Thus there is the generalized symmetry

$$s_{(+,-,+)} := \begin{pmatrix} -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

Since $(\mathfrak{a}_{u_0})_+ \neq 0$ is contained in $\mathfrak{p}_{fix}^{s_{(+,-,+)}}$, there is infinitely many automorphisms of type $\Sigma_{(+,-,+)}$.

But why we did not find any generalized symmetry of type $\Sigma_{(-,-,-)}$ although they are indicated in the Table 5? The reason is that $\lambda_{(-,-,-)}$ induces an outer automorphism of $Sp(8, \mathbb{R})$. Thus we have to consider $G' := \{B \in Gl(8, \mathbb{R}) : B\mathbb{J}B^T = \pm\mathbb{J}\}$, where \mathbb{J} gives the Lie group $Sp(8, \mathbb{R})$. It is not hard to check that the short exact sequence $Sp(8, \mathbb{R}) \rightarrow G' \rightarrow G'/Sp(8, \mathbb{R}) = \mathbb{Z}_2$ splits according to the choice of the block matrix $\begin{pmatrix} -\text{id} & 0 \\ 0 & \text{id} \end{pmatrix} \in G'/Sp(8, \mathbb{R})$. If we repeat the construction for

the type $(G', P'_{1,2,4})$, then we get the following additional generalized symmetries:

$$s_{(-,-,-)} := \begin{pmatrix} -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

and $s_{(-,+,-)} = s_{(-,-,-)} \circ s_{(+,-,+)}$. Since $(\mathfrak{a}_{u_0})_+ \neq 0$, there is infinitely many of them at any point.

Example 6 (Type $(\mathfrak{sl}(5, \mathbb{R}), \mathfrak{p}_{2,3})$ with curvature $(2, 1)$ or $(3, 2)$). (1) In this example, we will discuss three cases for the pair $(\mathfrak{sl}(5, \mathbb{R}), \mathfrak{p}_{2,3})$ with

$$\mathfrak{g}_- = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ x_1 & x_2 & 0 & 0 & 0 \\ x_4 & x_5 & x_3 & 0 & 0 \\ x_7 & x_8 & x_6 & 0 & 0 \end{pmatrix}.$$

- (2) We consider the harmonic curvatures $(2, 1)$, $(3, 2)$ or both of them at once.
 (3) The curvature in the case $\mu = (2, 1)$ is given by

$$\kappa(u_0)((x_1, \dots, x_8), (x'_1, \dots, x'_8)) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & x_1 x'_2 - x_2 x'_1 & 0 & 0 \end{pmatrix},$$

the curvature in the case $\mu = (3, 2)$ is given by

$$\kappa(u_0)((x_1, \dots, x_8), (x'_1, \dots, x'_8)) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ x_3 x'_5 - x_5 x'_3 & 0 & 0 & 0 & 0 \end{pmatrix},$$

and the curvature in the remaining case is the sum of the above.

- (4) In the case $\mu = (2, 1)$, $A_{u_0} = (A_{u_0})_0$ and is given by

$$\begin{pmatrix} x_9 & x_{10} & 0 & 0 & 0 \\ x_{11} & x_{12} & 0 & 0 & 0 \\ 0 & 0 & x_{13} & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{x_{13}} & 0 \\ 0 & 0 & 0 & x_{14} & \frac{x_{13}^3}{x_9 x_{12} - x_{10} x_{11}} \end{pmatrix}.$$

In the case $\mu = (3, 2)$, we compute

$$(A_{u_0})_0 = \begin{pmatrix} \frac{1}{x_{10}^3 x_{11}^3 x_{12}^3} & 0 & 0 & 0 & 0 \\ x_9 & x_{10} & 0 & 0 & 0 \\ 0 & 0 & x_{11} & 0 & 0 \\ 0 & 0 & 0 & x_{10}^2 x_{11}^2 x_{12}^2 & 0 \\ 0 & 0 & 0 & x_{13} & x_{12} \end{pmatrix},$$

and

$$(\mathbf{a}_{u_0})_+ = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & x_{14} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

Finally, in the case of both curvatures together, A_{u_0} is the intersection of both previous results, i.e.,

$$A_{u_0} = (A_{u_0})_0 = \begin{pmatrix} x_{11}^6 & 0 & 0 & 0 & 0 \\ x_9 & x_{10} & 0 & 0 & 0 \\ 0 & 0 & x_{11} & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{x_{11}^4} & 0 \\ 0 & 0 & 0 & x_{12} & \frac{1}{x_{10}x_{11}^3} \end{pmatrix}.$$

(5) In the case $\mu = (2, 1)$, elements of $Z(G_0) \cap A_{u_0}$ are of the form

$$s_{(a^{-5}, a^{-10})} = \begin{pmatrix} a^7 & 0 & 0 & 0 & 0 \\ 0 & a^7 & 0 & 0 & 0 \\ 0 & 0 & a^2 & 0 & 0 \\ 0 & 0 & 0 & a^{-8} & 0 \\ 0 & 0 & 0 & 0 & a^{-8} \end{pmatrix}$$

for $a \in \mathbb{R}$, and $s_{(-, +)}$ for $a = -1$ is the unique $s_{(-, +)}$ -symmetry.

In the case $\mu = (3, 2)$, elements of $Z(G_0) \cap A_{u_0}$ are of the form

$$s_{(a^5, a^4)} = \begin{pmatrix} a^{-3} & 0 & 0 & 0 & 0 \\ 0 & a^{-3} & 0 & 0 & 0 \\ 0 & 0 & a^2 & 0 & 0 \\ 0 & 0 & 0 & a^2 & 0 \\ 0 & 0 & 0 & 0 & a^2 \end{pmatrix}$$

for $a \in \mathbb{R}$, and $s_{(-, +)}$ for $a = -1$ is $s_{(-, +)}$ -symmetry and there is infinitely many of them.

In the case of both curvatures, $Z(G_0) \cap A_{u_0}$ is generated by $s_{(-, +)}$, which is the unique $s_{(-, +)}$ -symmetry.

7. TABLES

We present the tables describing the possibilities for the existence of non-trivial homogeneous parabolic geometries with generalized symmetries discussed in the Section 5. We will present here the complete list of the cases with \mathfrak{g} simple. Here:

- \mathfrak{g} ... simple Lie algebras and/or their real forms, if there are any restrictions on parameters, we will write them behind or below.
- Ξ ... the set of roots, which gives the parabolic subalgebra \mathfrak{p} of \mathfrak{g} , if there are any restrictions on parameters, we will write them below and we will omit the restrictions in the cases $\mathfrak{g} = \mathfrak{sl}(n+1, \mathbb{H}), \mathfrak{sp}(p, q)$, where the elements of Ξ are even.
- κ_H ... components of the harmonic curvature of the parabolic geometry associated to the pair $(\mathfrak{g}, \mathfrak{p})$, we indicate by ' that the corresponding root is in the other copy of \mathfrak{g} in the complexification of complex Lie algebra \mathfrak{g} .
- γ ... simple restricted roots that satisfy the condition (3) from the Theorem 4.3 for the given components of the harmonic curvature, if there are any restrictions on parameters, we will write them behind or below.

J ... all the endomorphisms J discussed in the Section 5 such that s_J acts trivially on the given component of the harmonic curvature, except the case that all complex structures J can be obtained by the composition of the given complex structure with the remaining entries. If the both $+$ and $-$ possibilities are available, we will simply write \pm on the corresponding position.

The source of the data for the pairs (\mathfrak{g}, Ξ) of simple parabolic geometries admitting a non-trivial curvature are the results in [1, 14]. The data about the components of the harmonic curvature and roots γ comes partly from the tables in [10], and partly are computed by the algorithm presented in [13]. Finally, using the results from the Section 5 and the explicit description of the components of the harmonic curvature, we compute, which elements s_J act trivially on the given components of the harmonic curvature.

We order the tables by the size of Ξ or number of non-trivial components of κ_H . We omit the cases which are equivalent to the other cases via outer automorphisms.

TABLE 2. The case $|\Xi| = 1$

\mathfrak{g}	Ξ	κ_H	γ	J
$\mathfrak{sl}(n+1, \{\mathbb{R}, \mathbb{C}\}), n > 2$	$\{1\}$	$(1, 2)$		$(-)$
$\mathfrak{sl}(n+1, \mathbb{C}), n > 1$	$\{1\}$	$(1, 1')$		$(-), (i)$
$\mathfrak{sl}(n+1, \{\mathbb{R}, \mathbb{C}, \mathbb{H}\}), n > 2$	$\{2\}$	$(2, 1)$		$(-)$
$\mathfrak{sl}(4, \{\mathbb{R}, \mathbb{C}, \mathbb{H}\})$	$\{2\}$	$(2, 1), (2, 3)$		$(-)$
$\mathfrak{sl}(n+1, \mathbb{C}), n > 2$	$\{1\}$	$(1, 2), (1, 1')$		$(-)$
$\mathfrak{so}(p, n-p), n > 6, p > 0$	$\{1\}$	$(1, 2)$		$(-)$
$\mathfrak{so}(n, \mathbb{C}), n > 6$	$\{1\}$	$(1, 2)$		$(-)$
$\mathfrak{sp}(2n, \{\mathbb{R}, \mathbb{C}\}), n > 2$	$\{1\}$	$(1, 2)$		$(-)$
$\mathfrak{sp}(2n, \mathbb{C}), n > 1$	$\{1\}$	$(1, 1')$		$(-), (i)$
$\mathfrak{sp}(2n, \{\mathbb{R}, \mathbb{C}\}), n > 2$	$\{2\}$	$(2, 1)$		$(-)$
$\mathfrak{sp}(p, n-p), n > 2, p > 0$	$\{2\}$	$(2, 1)$		$(-)$
$\mathfrak{sp}(2n, \mathbb{C}), n > 2$	$\{1\}$	$(1, 2), (1, 1')$		$(-)$
$\mathfrak{g}_2(2)$	$\{1\}$	$(1, 2)$		$(-)$
$\mathfrak{g}_2(\mathbb{C})$	$\{1\}$	$(1, 2)$		$(-), (i)$

TABLE 3. The case $|\Xi| = 2$, one component of κ_H

\mathfrak{g}	Ξ	κ_H	γ	J
$\mathfrak{su}(1, 2)$	$\{1, 2\}$	$(1, 2)$		$(-, -)$
$\mathfrak{sl}(n+1, \{\mathbb{R}, \mathbb{C}\}), n > 1$	$\{1, 2\}$	$(1, 2)$		(\pm, \pm)
$\mathfrak{sl}(n+1, \{\mathbb{R}, \mathbb{C}\}), n > 2$	$\{1, 2\}$	$(2, 1)$	α_1	$(+, -)$
$\mathfrak{sl}(n+1, \{\mathbb{R}, \mathbb{C}\}), n > 2$	$\{1, 3\}$	$(1, 2)$		$(-, +)$
$\mathfrak{sl}(n+1, \mathbb{R}), n > 2$	$\{1, n\}$	$(1, n)$		$(-, -)$
$\mathfrak{sl}(n+1, \mathbb{C}), n > 1$	$\{p, p+1\}$	$((p+1)', p')$	α_p	$(-, +)$
$\mathfrak{sl}(n+1, \mathbb{C}), n > 1$	$\{1, p\}$	$(1, p')$		$(+, -)$
$\mathfrak{su}(p, n-p), n > 3,$ $p > 0$	$\{1, n\}$	$(1, n)$		$(-, -),$ $(i, -i)$
$\mathfrak{sl}(n+1, \{\mathbb{R}, \mathbb{C}\})$ $n > 2$	$\{1, p\}$ $p > 3$	$(1, 2)$	α_p $n-1 > p$	$(-, +)$
$\mathfrak{sl}(n+1, \{\mathbb{R}, \mathbb{C}\})$ $n > 3$	$\{1, p\}$ $n > p > 2$	$(1, p)$		$(+, -)$
$\mathfrak{sl}(n+1, \mathbb{C})$ $n > 2$	$\{1, n\}$	$(1, n)$		$(-, -),$ $(i, -i)$
$\mathfrak{sl}(n+1, \{\mathbb{R}, \mathbb{C}, \mathbb{H}\})$ $n > 2$	$\{2, p\}$ $p > 2$	$(2, 1)$	α_p $n > p > 2$	$(-, +)$
$\mathfrak{sl}(n+1, \{\mathbb{R}, \mathbb{C}\})$ $n > 2$	$\{p, p+1\}$ $p > 1$	$(p+1, p)$	α_p $n > p+1$	$(-, +)$
$\mathfrak{sl}(n+1, \mathbb{C})$ $n > 1$	$\{1, p\}$	$(1, 1')$	α_p $n > p > 2$	$(-, +)$
$\mathfrak{so}(3, 4), \mathfrak{so}(7, \mathbb{C})$	$\{1, 3\}$	$(3, 2)$		$(-, -)$
$\mathfrak{so}(p, n-p), n > 4, p > 2$	$\{1, 2\}$	$(1, 2)$	$\alpha_2, n > 6$	(\pm, \pm)
$\mathfrak{so}(n, \mathbb{C}), n > 4$	$\{1, 2\}$	$(1, 2)$	$\alpha_2, n > 6$	(\pm, \pm)
$\mathfrak{so}(p, n-p), n > 6, p > 1$	$\{1, 2\}$	$(2, 1)$		$(-, +)$
$\mathfrak{so}(n, \mathbb{C}), n > 6$	$\{1, 2\}$	$(2, 1)$		$(-, +)$
$\mathfrak{so}(p, n-p), n > 6, p > 2$	$\{2, 3\}$	$(3, 2)$	$\alpha_2, n > 7$	$(-, +)$
$\mathfrak{so}(n, \mathbb{C}), n > 6$	$\{2, 3\}$	$(3, 2)$	$\alpha_2, n > 7$	$(-, +)$
$\mathfrak{so}(n, n), \mathfrak{so}(2n, \mathbb{C}), n > 3$	$\{1, n\}$	$(1, 2)$	$\alpha_n, n > 4$	$(-, +)$
$\mathfrak{sp}(2n, \{\mathbb{R}, \mathbb{C}\}), n > 1$	$\{1, 2\}$	$(1, 2)$		$(-, +)$
$\mathfrak{sp}(2n, \{\mathbb{R}, \mathbb{C}\}), n > 2$	$\{1, 2\}$	$(2, 1)$	α_1	$(+, -)$
$\mathfrak{sp}(2n, \{\mathbb{R}, \mathbb{C}\}), n > 2$	$\{1, n\}$	$(1, 2)$	$\alpha_n, n > 3$	$(-, +)$
$\mathfrak{sp}(2n, \{\mathbb{R}, \mathbb{C}\}), n > 2$	$\{1, n\}$	$(1, n)$		$(+, -)$
$\mathfrak{sp}(2n, \{\mathbb{R}, \mathbb{C}\}), n > 2$	$\{2, n\}$	$(2, 1)$	$\alpha_n, n > 3$	$(-, +)$
$\mathfrak{sp}(2n, \{\mathbb{R}, \mathbb{C}\}), n > 2$	$\{n, n-1\}$	$(n-1, n)$		$(+, -)$
$\mathfrak{sp}(n, n), n > 1$	$\{2, 2n\}$	$(2, 1)$	α_{2n}	$(-, +)$
$\mathfrak{sp}(2n, \mathbb{C}), n > 1$	$\{1, n\}$	$(1, 1')$	α_n	$(-, +)$
$\mathfrak{sp}(2n, \mathbb{C}), n > 1$	$\{1, n\}$	$(1, n')$		$(+, -)$
$\mathfrak{sp}(6, \mathbb{C})$	$\{2, 3\}$	$(2', 3')$		$(+, -)$
$\mathfrak{g}_2(2)$	$\{1, 2\}$	$(1, 2)$		(\pm, \pm)
$\mathfrak{g}_2(\mathbb{C})$	$\{1, 2\}$	$(1, 2)$		$(\pm, \pm), (i, i)$

TABLE 4. The case $|\Xi| = 2$, more component of κ_H

\mathfrak{g}	Ξ	κ_H	γ	J
$\mathfrak{sl}(3, \{\mathbb{R}, \mathbb{C}\})$	$\{1, 2\}$	$(1, 2), (2, 1)$		$(-, -)$
$\mathfrak{sl}(3, \mathbb{C})$	$\{1, 2\}$	$(1', 2'), (1, 2'), (2, 2')$		$(+, -)$
$\mathfrak{sl}(4, \mathbb{R})$	$\{1, 2\}$	$(2, 3), (1, 2), (2, 1)$		$(+, -)$
$\mathfrak{sl}(4, \mathbb{C})$	$\{1, 3\}$	$(1, 2), (1, 1'), (3, 1')$		$(-, +)$
$\mathfrak{sl}(4, \mathbb{C})$	$\{1, 3\}$	$(3, 2), (1, 3'), (3, 3')$		$(+, -)$
$\mathfrak{sl}(n+1, \mathbb{R}), n > 2$	$\{1, 2\}$	$(1, 2), (2, 1)$		$(+, -)$
$\mathfrak{sl}(n+1, \mathbb{C}), n > 3$	$\{1, 2\}$	$(1, 2), (2', 1'), (1, 1')$		$(-, +)$
$\mathfrak{sl}(n+1, \mathbb{C}), n > 3$	$\{1, 3\}$	$(1, 2), (1, 1')$		$(-, +)$
$\mathfrak{sl}(n+1, \mathbb{C}), n > 3$	$\{1, n\}$	$(1, 2), (1, 1'), (n, 1')$		$(-, +)$
$\mathfrak{sl}(n+1, \mathbb{R}), n > 2$	$\{2, n\}$	$(2, n), (2, 1)$		$(-, +)$
$\mathfrak{sl}(n+1, \mathbb{C}), n > 3$	$\{2, n\}$	$(2, n), (2, 1), (n, 2')$		$(-, +)$
$\mathfrak{sl}(n+1, \mathbb{C})$ $n > 3$	$\{1, p\}$ $p > 2$	$(1, p), (1, p')$		$(+, -)$
$\mathfrak{sl}(4, \mathbb{C})$	$\{1, 2\}$	$(2, 3), (1, 2), (2, 1),$ $(1, 2'), (1', 2')$		$(+, -)$
$\mathfrak{sl}(n+1, \mathbb{C})$ $n > 3$	$\{1, 2\}$	$(1, 2), (2, 1),$ $(1, 2'), (1', 2')$		$(+, -)$
$\mathfrak{sl}(n+1, \mathbb{C})$ $n > 3$	$\{1, p\}$ $p > 3$	$(1, 2), (1, 1')$	α_p	$(-, +)$
$\mathfrak{sl}(n+1, \mathbb{C})$ $n > 3$	$\{p, p+1\}$ $p > 2$	$(p+1, p), ((p+1)', p')$	$n-1 > p$ α_p	$(-, +)$
$\mathfrak{so}(p, n-p)$ $n > 4, p > 1$	$\{1, 2\}$	$(1, 2), (2, 1)$		$(-, +)$
$\mathfrak{so}(n, \mathbb{C}), n > 4$	$\{1, 2\}$	$(1, 2), (2, 1)$		$(-, +)$
$\mathfrak{sp}(4, \mathbb{C})$	$\{1, 2\}$	$(1, 2), (2, 1),$ $(1', 2'), (1, 2')$		$(-, +),$ $(+, -)$
$\mathfrak{sp}(2n, \mathbb{C})$ $n > 2$	$\{1, n\}$	$(1, 2), (1, 1')$	α_n	$(-, +)$
$\mathfrak{sp}(6, \mathbb{C})$	$\{2, 3\}$	$(2, 3), (2', 3')$		$(-, +)$
$\mathfrak{sp}(2n, \mathbb{C}), n > 2$	$\{1, n\}$	$(1, n), (1, n')$	$n > 3$	$(+, -)$

TABLE 5. The case $|\Xi| = 3, 4$, one component of κ_H

\mathfrak{g}	Ξ	κ_H	γ	J
$\mathfrak{sl}(n+1, \{\mathbb{R}, \mathbb{C}\})$	$\{1, 2, p\}$	$(1, 2)$	α_p	$(-, -, +),$
	$n > p$		$n > p > 3$	$(+, -, +), (-, +, +)$
$\mathfrak{sl}(n+1, \mathbb{R})$	$\{1, 2, p\}$	$(2, 1)$	α_1, α_p	$(-, -, -),$
			$n > p > 3$	$(-, +, -), (+, -, +)$
$\mathfrak{sl}(n+1, \mathbb{C})$	$\{1, 2, p\}$	$(2, 1)$	α_1, α_p	$(-, -, -), (-, +, -),$
			$n > p > 3$	$(+, -, +), (-i, i, i)$
$\mathfrak{sl}(n+1, \{\mathbb{R}, \mathbb{C}\})$	$\{1, p, n\}$	$(1, n)$	α_p	$(-, -, +),$
			$p > 2$	$(+, -, -), (-, +, -)$
$\mathfrak{su}(2, 2)$	$\{1, 2, 3\}$	$(2, 1)$		$(-, -, -),$
				$(-, +, -), (+, -, +)$
$\mathfrak{su}(n, n)$	$\{1, n,$ $2n-1\}$	$(1,$ $2n-1)$	α_n	$(-, +, -)$
			$n > 2$	
$\mathfrak{su}(2, 2)$	$\{1, 2, 3\}$	$(1, 2)$		$(+, -, +)$
$\mathfrak{so}(2n, \mathbb{C})$	$\{1, 2, n\}$	$(1, 2)$	α_2, α_n	$(-, -, +),$
$\mathfrak{so}(n, n), n > 3$			$n > 4$	$(+, -, +), (-, +, +)$
$\mathfrak{so}(7, \mathbb{C}),$ $\mathfrak{so}(3, 4)$	$\{1, 2, 3\}$	$(3, 2)$		$(-, -, -),$
				$(-, +, -), (+, -, +)$
$\mathfrak{sp}(2n, \{\mathbb{R}, \mathbb{C}\})$	$\{1, 2, p\}$	$(2, 1)$	α_1, α_p	$(+, -, -),$
	$p < n$			$(+, +, -), (+, -, +)$
$\mathfrak{sp}(2n, \{\mathbb{R}, \mathbb{C}\})$	$\{1, 2, n\}$	$(2, 1)$	α_1, α_n	$(-, -, -),$
			$n > 3$	$(-, +, -), (+, -, +)$
$\mathfrak{sl}(n+1, \{\mathbb{R}, \mathbb{C}\})$	$\{1, 2, p, q\}$	$(2, 1)$	$\alpha_1, \alpha_p, \alpha_q$	$(-, \pm, -, +),$
			$n > 4, p > 3,$ $p \neq n, q \neq n$	$(+, \pm, -, -),$ $(-, \pm, +, -)$
$\mathfrak{su}(n, n+1)$	$\{1, 2,$ $2n-1, 2n\}$	$(2, 1)$		$(-, +, +, -)$

 TABLE 6. The case $|\Xi| = 3, 4$, more component of κ_H

\mathfrak{g}	Ξ	κ_H	γ	J
$\mathfrak{sl}(4, \{\mathbb{R}, \mathbb{C}\})$	$\{1, 2, 3\}$	$(2, 1), (2, 3),$ $(1, 3)$		$(-, +, -)$
$\mathfrak{sl}(4, \{\mathbb{R}, \mathbb{C}\})$	$\{1, 2, 3\}$	$(2, 1), (2, 3),$ $(1, 2), (3, 2)$		$(+, -, +)$
$\mathfrak{sl}(n+1, \{\mathbb{R}, \mathbb{C}\})$	$\{1, 2, p\}$	$(1, 2), (2, 1)$	α_p	$(+, -, +)$
			$n > p > 3$	
$\mathfrak{sl}(4, \{\mathbb{R}, \mathbb{C}\})$	$\{1, 2, 3\}$	$(2, 1), (2, 3)$		$(-, -, -)$
$\mathfrak{su}(2, 2)$	$\{1, 2, 3\}$	$(1, 3), (2, 1)$		$(-, +, -)$
$\mathfrak{su}(2, 2)$	$\{1, 2, 3\}$	$(1, 2), (2, 1)$		$(+, -, +)$
$\mathfrak{sl}(n+1, \{\mathbb{R}, \mathbb{C}\})$	$\{1, 2, n\}$	$(1, 2), (1, n)$		$(-, -, +)$
$\mathfrak{sl}(n+1, \mathbb{R})$	$\{1, 2, n\}$	$(1, n), (2, 1)$		$(-, +, -)$
$\mathfrak{so}(4, 4), \mathfrak{so}(8, \mathbb{C})$	$\{1, 2, 4\}$	$(1, 2), (4, 2)$	α_2	$(+, -, +)$
$\mathfrak{sl}(n+1, \{\mathbb{R}, \mathbb{C}\})$	$\{1, 2, n-1, n\}$	$(2, 1), (n-1, n)$		$(-, -, -, +),$ $(+, -, -, -),$ $(-, +, +, -)$

REFERENCES

- [1] Čap A., and J. Slovák, “Parabolic Geometries I: Background and General Theory”, Amer. Math. Soc., 2009.
- [2] Čap A., and J. Slovák, V. Souček, *Bernstein-Gelfand-Gelfand Sequences*, Ann. of Math. **154**, (2001), 97-113.
- [3] Čap A., and K. Melnick, *Essential Killing fields of parabolic geometries*, Indiana Univ. Math. J. **62**, (2013), 1917-1953.
- [4] Gregorovič J., *General construction of symmetric parabolic geometries*, Differential Geometry and its Applications **30**, (2012), 450-476
- [5] Gregorovič J., “Geometric structures invariant to symmetries”, Masaryk University, 2012.
- [6] Gregorovič J., *Classification of invariant AHS-structures on semisimple locally symmetric spaces*, Central European journal of mathematics **11**, (2013), 2062-2075.
- [7] Hammerl M., *Homogeneous Cartan geometries*, Archivum Mathematicum (Brno) **43**, (2007), 431-442.
- [8] Knapp A.W., “Lie Groups Beyond an Introduction”, Birkhäuser, 1996.
- [9] Kolář I., and P.W. Michor, J. Slovák, “Natural operations in differential geometry”, Springer-Verlag, Berlin-Heidelberg-New York, 1993.
- [10] Kruglikov B., and D. The, *The gap phenomena in parabolic geometries*, arXiv:1303.1307v4
- [11] Loos O., *Spiegelungsraume und homogene symmetrische Raume*, Math. Z. **99**, (1967), 141-170.
- [12] Sharpe R. W., “Differential geometry: Cartan’s generalization of Klein’s Erlangen program”, Springer-Verlag, 1997.
- [13] Šilhan J., *A real analog of Kostant’s version of the Bott-Borel-Weil theorem*, Journal of Lie Theory **14**, (2004), 481-499
Algorithm available on: <http://bart.math.muni.cz/~silhan/lie/>
- [14] Yamaguchi K., *Differential systems associated with simple graded Lie algebras*, Advanced Studies in Pure Mathematics **22**, Progress in Differential Geometry, (1993), 413-494.
- [15] Zalabová L., *Symmetries of Parabolic Geometries*, Differential Geometry and its Applications **27**, (2009), 605-622.
- [16] Zalabová L., *Symmetries of Parabolic Contact Structures*, Journal of Geometry and Physics **60**, (2010), 1698-1709.

JAN GREGORVIČ, DEPARTMENT OF MATHEMATICS, AARHUS UNIVERSITY, NY MUNKEGADE 118, AARHUS C 8000, DENMARK, AND, DEPARTMENT OF MATHEMATICS, AND STATISTIC, FACULTY OF SCIENCES, MASARYK UNIVERSITY, KOTLÁŘSKÁ 2, BRNO 611 37, CZECH REPUBLIC, JAN.GREGOROVIC@SEZNAM.CZ

LENKA ZALABOVÁ, INSTITUTE OF MATHEMATICS AND BIOMATHEMATICS, FACULTY OF SCIENCE, UNIVERSITY OF SOUTH BOHEMIA, BRANIŠOVSKÁ 31, ČESKÉ BUDĚJOVICE 370 05, CZECH REPUBLIC, LZALABOVA@GMAIL.COM